MAEDICA – a Journal of Clinical Medicine 2016; 11(4): 286-295

ORIGINAL PAPER

Nutritional Knowledge, Attitude and Practice of North West Households in Iran: Is Knowledge likely to Become Practice?

Sakineh Nouri SAEIDLOU^a, Fariba BABAEI^b, Parvin AYREMLOU^a

^a Food and Beverage Safety Research Center, Urmia University of Medical Science, Urmia, Iran

^b Urmia University of Medical Science, Urmia, Iran

-ABSTRACT-

Background: Nutritional factors have an important role in the incidence of non-communicable diseases and they are strong predictors of cardiovascular risk among adolescents. This study was conducted to assess the Knowledge, Attitude and Practice (KAP) of urban and rural households towards the principles of nutrition in West Azerbaijan Province.

Design and methods: This cross sectional population based study was conducted among 455 urban and rural households. The sampling method at households' level in each area was the single stage cluster sampling. Data were collected by a structured questionnaire and through the interview with the eligible subject in each household. Analyses were performed using SPSS 20 statistical software. For qualitative data, results were presented as frequency and percentage.

Results: Out of a total of 455 households that were selected as overall sample size, 272 (59.8%) were in urban areas and 183 (40.2%) in rural areas. More than 50% of the households in both urban and rural areas were aware of food groups. More than 40% of the households in both urban and rural areas knew about the roles of food groups and the level of knowledge most frequently encountered was about the role of milk and dairy group in urban areas (88.6%). Vitamin intake associated with the role of fruits (68%) and vegetables (62.5%) had the highest frequency. Most of the respondents declared that they consumed certain foods for their effects related to health improvement and disease prevention. The results showed that 45.2-99.8% of them had favorable attitudes. Most of the households consumed red meat, poultry, egg and legumes weekly, whereas fish was eaten rarely. Fruits, vegetables, sugar and dairy were consumed daily in most of the households.

Conclusion: Higher nutrition knowledge is associated with better practice and diet quality. Our findings suggest that nutrition education should be integrated in a household training program.

Keywords: Nutrition, Knowledge, Attitude, Practice, Household, Iran

Address for correspondence:

Parvin Ayremlou

Article received on the 24rd of October 2016. Article accepted on the 14rd of December 2016.

Postal address: Food and Beverages Safety Research Center, Gholipour street, Jaddeh Salmas, Urmia, Iran

Phone number: 009844-33457278, Fax: 00984433457277;

E-mail: p.ayremlou@gmail.com, saeedlou2003@yahoo.com, babaei707@yahoo.com, babaei707@yabaei707000?wyabaei707@yabaei707@y

INTRODUCTION

he World Health Organization has stated that nutrition is an input to and foundation for health and development. Proper nutrition makes people stronger and more productive (1). Healthy eating habits lead to a stronger immune system, less illness, and better health. Proper and healthy nutrition is a key to a better quality of life (2-6). Epidemiological studies have shown that changes in lifestyle during recent years, especially changes in nutritional habits, diet, types of food, and cooking time may be responsible for increasing rates of non-communicable diseases (7-10). Nutritional factors have an important role in the incidence of non-communicable diseases and they are strong predictors of cardiovascular risk among adolescents (11-13). Epidemiological evidence shows that there is an increased incidence of diseases such as Cardiovascular diseases, obesity, high blood pressure and cancer, which can be attributed to changes in both lifestyle and nutritional habits (14-16).

One of the main causes of nutritional problems, the lack of nutrition knowledge, results in poor practice, which causes serious harms such as malnutrition and various non-communicable diseases (17, 18). KAP (Knowledge, Attitudes and Practices) studies emerged in the 1950s from the need to measure opposition to family planning services (19). Since then, they have been used extensively in family planning and population studies to evaluate and guide existing programmes and their use has been extended to other areas of health, including nutrition (20). Nutrition-related KAP studies assess and explore peoples' KAP relating to nutrition, diet, foods as well as closely related hygiene and health issues (21, 22). KAP studies have been used for two main purposes: (I) to collect key information during a situation analysis, which can then feed into the design of nutrition interventions; and (II) to evaluate nutrition education interventions. In the context of nutrition-related projects or programmes, a situation analysis describes the type and magnitude of nutrition issues and identifies possible causes of the commonly observed nutritional problems. The findings of a situation analysis will help in planning a nutrition intervention aimed at alleviating the identified nutritional problems. KAP studies can contribute to a situation analysis by helping to determine the existing knowledge, attitudes and practices relating to nutrition, which identifies nutrition education priorities (23).

Iran, like many developing countries, is facing an increase in the prevalence of risk factors for non-communicable diseases due to changes in lifestyle, especially nutritional habits (9, 24, 25). Several studies indicate the relationship between nutrition and various diseases that may be due to poor knowledge and practices in nutritional habits (26-28). Some studies conducted in this region show that malnutrition and obesity has increased due unhealthy dietary habits (29-31). Therefore, using a dietary pattern method which considers a more comprehensive overview of the diet could provide more interpretable findings than studying single nutrients or foods. Awareness concerning the level of consumers' nutritional knowledge is useful for promoting dietary habits and teaching good eating habits, and proper nutritional behaviors lead to reduced disease and treatment costs (32-36). Given the high prevalence of non-communicable diseases and their risk factors, nutritional intervention is necessary to prevent these diseases (37). This study provides baseline data on the knowledge, attitude and practice of urban and rural households in North West (West Azerbaijan Province) of Iran for planning nutritional intervention programmes. 🗖

MATERIAL AND METHOD

This cross sectional population based study was conducted to assess the Knowledge, Attitude and Practice (KAP) of urban and rural households towards the principles of nutrition in West Azerbaijan Province. We considered a mother or any member of the household who was above 15 years old and responsible for cooking meals in the family as index case for data gathering.

Subjects and sampling

The sampling method at households' level in each area was the single stage cluster sampling with equal size clusters (8 people in each cluster). Regarding the design effect of 1.5, the total sample size was estimated to be 456 (34 cluster in urban regions and 23 cluster in rural regions),

Variables		Urban (%)	Rural (%)	Total (%)
Sex	Male Female	13 (4.8) 259 (95.2)	3 (1.6) 180 (98.4)	16 (3.5) 439 (96.5)
Age group	<20 years old 20-39 years old 40-64 years old > 65 years old	3 (1.1) 122 (44.9) 128 (47.1) 19 (7)	6 (3.3) 99 (54.1) 73 (39.9) 5 (2.7)	9 (2) 221 (48.6) 201 (44.2) 24 (5.3)
Education	Illiterate Elementary school Middle school High school University	54 (19.9) 55 (20.2) 38 (14) 82 (30.1) 43 (15.8)	85 (46.7) 58 (31.9) 14 (7.7) 18 (9.9) 7 (3.8)	139 (30.6) 113 (24.9) 52 (11.5) 100 (22) 50 (11)

TABLE 1. Demographic characteristics of respondents in urban and rural households

but finally, 455 people took part in this study. To identify clusters, we obtained the list of all households in each population (urban and rural) from deputy of health based on last health census. For finding first household of each cluster, we used systematic random sampling. We provided the cumulative list of households in each area. The sampling interval was calculated (number of clusters in each area divided by the total number of households in the area). The first household of the first cluster was identified by randomly selecting a number between 1 and a figure lower than or equal to the sampling interval. The first household of the second cluster was located by adding the sampling interval to the random

Questions		Urban (%)	Rural (%)	Total (%)
1-Recognition of food	Bread, Grains and Rice	212 (77.9)	130 (71)	342 (75.2)
groups	Meat, Legume and Egg	227 (83.5)	130 (71)	357 (78.5)
	Milk and dairy group	170 (62.5)	91 (50)	261 (57.5)
	Fruits	157 (57.7)	80 (43.7)	237 (52.1)
	Vegetables	184 (67.6)	98 (53.6)	282 (62)
2-Roles of food groups	Bread, Grains and Rice	111 (40.8)	95 (51.9)	206 (45.3)
	Meat, Legume, Egg	140 (51.7)	56 (30.6)	196 (43.2)
	Milk and dairy group	240 (88.6)	109 (59.6)	349 (76.9)
3-Roles of fruits	Vitamin intake	223 (82.3)	85 (46.7)	308 (68)
	Mineral intake	56 (20.7)	56 (30.6)	112 (24.7)
	Dietary fiber intake	42 (15.5)	22 (12)	64 (14.1)
4-Roles of vegetables	Vitamin intake	204 (75.3)	92 (50.3)	296 (65.2)
	Mineral intake	78 (58.8)	51 (27.9)	129 (28.4)
FD (() ()	Dietary fiber intake	63 (23.3)	45 (24.6)	108 (23.8)
5-Reason of food eating	Growth	102 (37.5)	61 (33.3)	163 (35.8)
	Energy intake	154 (56.8)	110 (60.4)	264 (58.3)
	Health and prevention of diseases	172 (63.2)	104 (56.8)	276 (60.4)
6-Effect of high	Overweight and obesity	115 (42.4)	50 (27.3)	165(36.3)
consumption of fast foods	Liver diseases	18(6.6)	11(6)	29(6.4)
	Candiana and an dianana	54 (19.9)	22(12)	76 (16.7) 81 (17.8)
	Cardiovascular diseases	58 (21.4) 20 (25.1)	23(12.6)	81(17.8)
7 Deles of fibers	Prevention of cancers	20(35.1)	7 (15.9)	27 (26.7)
7-Roles of libers	Help to howel movements	10(20.1)	7(10.9) 17(45.0)	23(24.3)
	Provention of overweight and obesity	30(00.7)	17(43.9) 11(28.5)	33(36.3)
8 Efforts of fizzy dripks	Overweight and obesity	23(39.7)	11(20.3)	34(33.4) 146(32.4)
bigh consumption	Ostooporosis	104(384)	47(20)	140(32.4) 153(33.7)
nightconsumption	Malnutrition	52(192)	25(137)	77(17)
	The loss of tooth enamel	28(10.4)	18 (9 9)	46(102)
9-Fat group	The liquid oil is the best oil for cooking	20(10.4) 219(80.8)	10(9.9) 124(67.8)	343(756)
y rut group	The liquid oil is the best oil for frying	220 (81 5)	121(07.0) 117(63.9)	337(744)
10-Harmful oils	Ghee	115 (42.9)	76 (41.5)	191(424)
10 114111141 0115	Solid	214 (79)	130(71)	344 (75.8)
11-Protein intake of	Sova	148 (54.6)	58 (31.7)	206 (45.4)
non-meat group	Legumes	145 (53.5)	76 (41.5)	221 (48.7)
	Egg	97 (35.8)	75 (41)	172 (37.9)
	Milk& dairy products	50 (18.5)	54 (29.5)	104 (22.9)

TABLE 2. Correct nutrition knowledge of urban and rural households in West Azarbijan Province

number. Non-Iranian households were excluded from the study. Also, households who were not present at the time of interview for three times were excluded from the survey.

Date collection

Data was collected by a structured guestionnaire and through an interview with the eligible subject in each household. Interviews were done by trained experts. The questionnaires focused on the basic principles of nutrition, food groups, and sources of nutrients, nutrition-related diseases, and nutritional requirement at different life stages. The questionnaires were standardized and pilot tested for validity and reliability. The study manual of operations was evaluated and revised in the pilot study. The knowledge status was assessed by 11 items and categories of response were she/he knows and she/he does not know. Attitude was measured by 19 items and response categories were from 1 totally agree to 5 totally disagree. Practice was assessed by 23 items and answers ranged from daily to never. All analyses were performed using SPSS 20 statistical software. For qualitative data, results were presented as frequency and percentage. \Box

RESULTS

A total of 455 households were selected as overall sample size, of which 272 (59.8%) were located in urban areas and 183 (40.2%) in rural areas. More than 95% of the respondents were women; 47.1% of them were in the age group of 40 to64 years in urban areas, and 54.1% in the age group of 20 to39 years in rural areas. The most frequent category of education reported by the respondents was high school (30.1%) in urban areas and illiterate (46.7%) in rural areas (Table 1).

Table 2 shows the knowledge level of urban and rural households toward nutrition. More than 50% of households in both urban and rural areas were aware of food groups, except for the frequency of the level of knowledge about fruits, which was 43.7% in rural areas. More than 40% of households in both urban and rural areas knew about the roles of food groups and the most frequently encountered level of knowledge was about role of milk and dairy group in urban areas (88.6%). Vitamin intake associated with

the role of fruits (68%) and vegetables (62.5%) had the highest frequency. Most of the respondents declared that they consumed certain foods for their effects regarding health improvement and disease prevention. 42.4% and 27.3% of the urban and rural households, respectively, knew that high consumption of fast foods was a main risk factor for overweight and obesity. Totally, 58.5% of the households were aware that the role of fibers in the body was useful for bowel movements. Awareness of osteoporosis as a health problem related to the high consumption of fizzy drinks had the highest frequency among respondents (33.7%). Most of the households in both urban and rural areas declared that liquid oil was the best for cooking and frying, while 75.8% of all respondents knew that solid oil was harmful. The best level of knowledge about protein intake of non-meat group was about soya (54.6%) and legumes (48.7%) in urban and rural areas, respectively.

Table 3 highlight the favorable attitudes of households toward basic principles of nutrition. The results showed that 45.2-99.8% of the families had favorable attitudes in all items, except that only 24.4% of the households believed that the nutritional value of meat and mushrooms was different. More than 90% of the households had favorable attitudes about (1) the importance of nutrition and diet in human health, (2) the necessity to pay more attention to the nutritional needs of children than to those of adults, (3) the necessity of a daily intake of vegetables or salad, (4) the priority of drinking buttermilk instead of soda, and (5) not using chips as a snack for children.

Table 4 shows the practice of consumption based on region. Findings indicated that most of the households have never consumed fast foods (Sausages (55.5%), Pizza (46.3%) and Sandwiches (54.8%). Among all households, the frequency of daily consumption of fruits and vegetables was 84.8% and 78.9%, respectively. Most of the households consumed fish, tons, fizzy drinks and industrial juice rarely in both urban and rural areas. Results showed that more than half of the households consumed buttermilk daily, while milk, yoghurt or cheese were eaten daily by most of the urban and rural families (urban households 89.3% and rural households 90.7%). Most of the households have never used snacks. The frequency of weekly egg intake was 62.4

Topic	Urban (%)	Rural (%)	Total (%)
1-The importance of nutrition and diet in human health	271 (100)	182 (99.5)	453 (99.8)
2-The necessity of same food intake in both sexes when a small	192 (70.8)	120 (65.6)	312 (68.7)
amount of food is in the households basket			
3-The necessity to pay more attention to the nutritional needs of	261 (96.3)	177 (96.7)	438 (96.5)
children than to those of adults			
4-The necessity of eating main meals in weight loss diets	157 (58.4)	97 (53.6)	254 (56.4)
5- The priority of traditional foods upon fast food	249 (91.9)	148 (80.9)	397 (87.4)
6-The observing fitness in girls at puberty with consume less food	160 (59)	109 (59.6)	269 (59.3)
7-The priority of using fruits instead of bread when she/he is hungry	105 (38.7)	100 (54.6)	205 (45.2)
8- The necessity to have a daily intake of vegetables or salad	261 (96.3)	175 (95.6)	436 (96)
9- Reducing vitamins in vegetables by freezing	231 (85.2)	121 (66.1)	352 (77.5)
10-The priority of fish consumption instead tons	198 (73.1)	99 (54.1)	297 (65.4)
11- The priority of drinking buttermilk instead soda	246 (91.8)	165 (90.7)	411 (91.3)
12- Using of liquid oils instead of solid oils	231 (85.2)	125 (68.3)	356 (78.4)
13-The harmful consumption of fizzy drinks	201 (74.4)	117 (63.9)	318 (70.2)
14- The priority of low-fat dairy products	194 (71.6)	94 (51.4)	288 (63.4)
15- The daily milk intake in all age groups	230 (85.2)	161 (88.5)	391 (86.5)
16- Not using chips as a snack for children	256 (94.5)	163 (89.1)	419 (92.3)
17- Difference in the nutritional value mushrooms and meat	65 (24.2)	45 (24.7)	110 (24.4)
18- The priority of boiled foods	234 (86.3)	161 (88)	395 (87)
19- Wholemeal bread has more nutritional value than white bread	139 (51.9)	79 (43.2)	218 (48.3)

TABLE 3. Favorite nutrition attitude of urban and rural households in West Azarbijan Province

and 53.6% in urban and rural households, respectively. Red meat, legumes and poultry were eaten weekly in most of the households (red met 68.7%, legumes 49.6% and poultry 80.6%). Butter and cream were eaten daily by most of the urban families (37.7%), whereas most of the rural families (38.7%) consumed it weekly. 84.4% of the urban families and 90.6% of the rural families consumed sugar daily.

Table 5 shows the nutrition practice of urban and rural households regarding the type of dairy, bread, and oil consumption. Findings indicated that in all households, the consumption frequency of traditional and pasteurized (low fat) dairy was 50.3%, 40.8% and 8.8%, respectively. In both urban and rural areas, most of the families consumed white bread. Among households, the consumption frequencies of liquid oils for cooking and frying oils for frying were higher comparing to other types of oils.

DISCUSSION

ncreasing epidemic of obesity and chronic diseases is associated to unhealthy dietary habits worldwide (38-42). Raising people's awareness for healthy dietary habits is a complex task and therefore, a better knowledge of these patterns is required. Studies that access and analyse people's nutrition-related knowledge, attitudes and practices (KAP) are a useful method for gaining such an insight into peoples' personal determinants of their dietary habits. They can thus provide valuable inputs for effective program and project planning.

The aim of this study was to assess knowledge, attitude and practice of urban and rural households toward the principles of nutrition. According to our findings, the urban and rural households' level of knowledge about recognition of food groups was acceptable. The results showed that most of the households were aware about the role of milk and dairy group, whereas the lowest level of knowledge frequency was for meat, legume and egg group. Most of the families declared that the importance of fruits and vegetables intake consists of providing vitamins, minerals and dietary fibers, respectively. Most of the households reported that they preferred to eat certain foods for their effects regarding health improvement and disease prevention. In both urban and rural areas, the households level of knowledge about the dairy group and the importance of fruits and vegetables in providing vitamins was good and acceptable, but it was weak about bread, grain and meat group as well as the role of fruits/vegetables in providing mineral and dietary fiber. Therefore, nutrition edu-

	Urban (%)	Rural (%)	Total (%)		
1-Sausage					
Daily	8 (3)	5 (2.7)	13 (2.9)		
Weekly	11 (4.1)	10 (5.5)	21 (4.6)		
Rarely	113 (41.7)	55 (30.1)	168 (37)		
Never	139 (51.3)	113 (61.7)	252 (55.5)		
2-Pizza					
Daily	12 (4.4)	9 (4.9)	21 (4.6)		
Weekly	11 (4.1)	5 (2.7)	16 (3.5)		
Karely	141(52) 107(20E)	66(36.1) 102(56.2)	207 (45.6)		
3-Sandwich	107 (39.3)	105 (50.5)	210 (40.3)		
Daily	7 (2 6)	8 (4 4)	15 (3.3)		
Weekly	27(10)	14 (7.7)	41 (9)		
Rarely	107 (39.5)	42 (23)	149 (32.8)		
Never	130 (48)	119 (65)	249 (54.8)		
4-Fruits					
Daily	233 (87.3)	131 (72.8)	385 (84.8)		
Weekly	29 (10.9)	38 (21.1)	49 (10.8)		
Rarely	3 (1.1)	7 (3.9)	5 (1.1)		
Never	2 (0.7)	4 (2.2)	15 (3.3)		
5-Vegetables	215(70.2)	142 (79.1)	259 (79 0)		
Dany	213(79.3)	143(78.1) 24(18.6)	338 (78.9) 81 (17.8)		
Rarely	$\frac{47}{5(18)}$	2(10.0)	7(15)		
Never	4(1.5)	4(22)	8 (1.8)		
6-Fish	1 (1.0)	- ()	0 (1.0)		
Daily	3 (1.1)	3 (1.6)	6 (1.3)		
Weekly	73 (26.9)	46 (25.1)	119 (26.2)		
Rarely	152 (56.1)	107 (58.5)	259 (57)		
Never	43 (15.9)	27 (14.8)	70 (15.4)		
7-Tons					
Daily	2 (0.7)	1 (0.5)	3 (0.7)		
Weekly	48 (17.7)	48 (26.2)	96 (21.1)		
Karely	148(54.6)	99 (54.1) 25 (10.1)	247 (54.4)		
8-Fizzy drinks	75 (20.9)	55 (19.1)	100 (23.0)		
Daily	11 (4.1)	17 (9.3)	28 (6.2)		
Weekly	53 (19.2)	36 (19.7)	88 (19.4)		
Rarely	155 (42.4)	89 (48.6)	244 (44.9)		
Never	93 (34.3)	41 (22.4)	134 (29.5)		
9-Industrial juice					
Daily	16 (5.9)	12 (6.6)	28 (6.2)		
Weekly	51 (18.8)	37 (20.2)	88 (19.4)		
Rarely	115 (42.4)	85 (46.4)	200 (44.1)		
10 Protterme ille	89 (32.8)	49 (26.8)	138 (30.4)		
10-Duttermik	148 (55.2)	126 (68 9)	247 (60.8)		
Wookly	96 (35.8)	42 (23)	138 (30.6)		
Rarely	19 (7.1)	9 (4 9)	28 (6.2)		
Never	5 (1.9)	6 (3.3)	11 (2.4)		
11-Milk, yoghurt or cheese					
Daily	242 (89.3)	166 (90.7)	408 (89.9)		
Weekly	22 (8.1)	13 (7.1)	35 (7.7)		
Rarely	6 (2.2)	3 (1.6)	9 (2)		
Never	1 (0.4)	1 (0.5)	2 (0.4)		

TABLE 4 (part one). Nutrition practice of urban and rural households in West Azarbijan Province

		Urban (%)	Rural (%)	Total (%)
12-Snacks				
	Daily	19 (7)	11 (6)	30 (3.6)
	Weekly	49 (18.1)	38 (20.8)	87 (19.2)
	Rarely	77 (28.4)	59 (35.2)	136 (30)
	Never	126 (46.5)	75 (41)	20 (44.3)
13-Chocolate	D 1	25 (0.4)		20 (0 7)
	Daily	25 (9.4)	14(7.7)	39 (8.7) 110 (26 F)
	VVeekiy Dorolu	71(20.7) 115(42.2)	48 (26.2)	119(20.5) 105(42.4)
	Novor	113(43.2) 55(20.7)	00(43.7)	195(43.4) 96(21.4)
14-Foos	INEVEL	55 (20.7)	41 (22.4)	90 (21.4)
11-1555	Daily	88 (32.5)	68 (37.2)	156 (34.4)
	Weekly	169 (62.4)	98 (53.6)	267 (58.8)
	Rarely	12 (4.4)	11 (6)	23 (5.1)
	Never	2 (0.7)	6 (3.3)	8 (1.8)
15-Red meet				
	Daily	34 (12.5)	18 (9.8)	52 (11.5)
	Weekly	187 (69)	125 (68.3)	312 (68.7)
	Rarely	45 (16.6)	35 (19.1)	80 (17.6)
4 6 7	Never	5 (1.8)	5 (2.7)	10 (2.2)
16-Legumes		110 (40 5)	00 (44.0)	000 (44.1)
	Daily	118 (43.5) 124 (40.4)	82 (44.8)	200(44.1)
	Paraly	134(49.4) 12(4.8)	91(49.7) 5 (2.7)	223 (49.6) 18 (4)
	Never	6(22)	5(2.7) 5(2.7)	10(4) 11(24)
17-Poultry	INCVCI	0 (2.2)	5 (2.7)	11 (2.1)
i, i outry	Daily	29 (10.7)	24 (13.1)	53 (11.7)
	Weekly	225 (83)	141 (77)	366 (80.6)
	Rarely	14 (5.2)	12 (6.6)	26 (5.7)
	Never	3 (1.1)	6 (3.3)	9 (2)
18-Butter and cream				
	Daily	71 (26.2)	69 (37.7)	140 (30.8)
	Weekly	105 (38.7)	58 (31.7)	163 (35.9)
	Rarely	65 (24)	41 (22.4)	106 (2.3)
10.0	Never	30 (11.1)	15 (8.2)	45 (9.9)
19-Sugar	D. 1	007 (04 4)	1(1(00))	201 (9(0)
	Daily	227 (84.4)	164 (90.6) 7 (2.0)	391 (86.9)
	Rarely	10(3.7) 15(5.6)	9 (5)	$\frac{17}{(3.8)}$
	Never	17 (6.3)	1 (0 5)	18(4)
		17 (0.0)	1 (0.0)	10(1)

TABLE 4 (part two). Nutrition practice of urban and rural households in West Azarbijan Province

cation is needed to get better knowledge of these aspects. Our finding showed that most of the households knew that fast food consumption could lead to overweight and obesity (36.3%), fibers were helpful for bowel movements (58.5%), fizzy drinks intake was a risk factor for osteoporosis (33.7%) and most of them (more than 70%) reported that liquid oil was the best oil

In a study by Alaunyte et al., the findings revealed an adequate nutritional knowledge (mean 72.82%) of athletes with the highest scores in dietary advice section (85.71%), followed by food groups (71.24%) and food choice (69.52%)(4). Fallah F et al. investigated the Effects of Nutrition Education on Levels of Nutritional Awareness of Pregnant Women in Western Iran. Their results indicated that, before the educational intervention, awareness levels were estimated to be weak (31%), moderate (66%) and good (3%). The corresponding rates after the educational intervention were weak (6%), moderate (63%) and good (31%). Pregnant women's awareness level about healthy nutrition was significantly increased from 3% before the intervention to 31% after the nutritional education

		Urban (%)	Rural (%)	Total (%)
1-Dairy				
, ,	Traditional	114 (42.2)	114 (62.3)	228 (50.3)
	Pasteurized (low fat)	127 (47)	58 (31.7)	185 (40.8)
	Pasteurized(high fat)	29 (10.7)	11 (6)	40 (8.8)
2-Bread				
	Lavash (white)	233 (86)	134 (73.6)	367 (81)
	Taftoon (white)	7 (2.6)	5 (2.7)	12 (2.6)
	Sangak (wholegrain)	26 (9.6)	18 (9.9)	44 (9.7)
	Barbary (wholegrain)	2 (0.7)	0 (0)	2 (0.4)
	Baget (white)	2 (0.7)	2 (1.1)	4 (0.9)
	Traditional (white)	1 (0.4)	23 (12.6)	24 (5.3)
3-Cooking oil				
	Solid oils	87 (32.2)	67 (37)	154 (34.2)
	Liquid oil	130 (48.3)	69 (38)	199 (44.3)
	Butter and others	52 (19.4)	45 (25)	97 (21.5)
4-Frying oil				
	Solid oils	37 (13.7)	36 (19.7)	73 (16.1)
	Liquid oil	25 (9.3)	29 (15.8)	54 (11.9)
	Frying oils	183 (67.5)	92 (50.3)	275 (60.6)
	Butter and others	26 (2.5)	25 (14.2)	52 (11.4)

TABLE 5. Nutrition practice of urban and rural households about type of dairy, bread, oil consumption in West Azarbijan Province

intervention (32). Sichert-Hellert *et al.* reported that in the total sample, more than 50% of the respondents were answered correctly and between 22% and 93% of participants answered the single questions correctly. Webb *et al.* reported that respondents gave correct answers to less than 50% of the survey questions. Participants scored below 50% for 10 of the 21 nutrition knowledge questions (43).

The current study showed that 45.2-99.8% of the families had favorable attitudes in all items, except that only 24.4% of the households believed that the nutritional value of meat and mushrooms was different. Webb et al. reported that the mean score for attitude towards nutrition for the entire sample was 41.69 ± 6.215 . The highest score was 55 (100%) and the lowest 19 (34.55%)(33). In a study by Sharma et al. when assessing nutrition attitudes, 54% of the teachers agreed with the statement that it was hard to know which nutrition information to believe and 9% agreed that their nutrition habits were healthy. Most teachers (78%) agreed that being overweight increases health risks, and 93% agreed that learning the relationship between health and nutrition was important (44).

In the present study, more than half of the households have never consumed sausages and sandwiches, while less than 5% of them eated

fast foods daily. Positive associations were found between fast food consumption and increased risk of abdominal obesity and hypertriglyceridemia. Sausage, French fries and pizza consumption was associated with a higher risk of incident metabolic syndrome (45). In both urban and rural areas, more than 70% of the household's fruits and vegetables intake was daily; therefore, nutritional knowledge was positively correlated to fruits and vegetables consumption by families. In Sharma's et al. study, nearly 1/4 of the teachers reported that they had not eaten fruits (26%) or vegetables (23%) the previous day, one of three (32%) consumed at least one fruit flavored beverage (e.g., fruit punch, sports drinks), and 44% had at least one soda on the previous day (44). The present study showed that the frequency of daily, weekly and seldom consumption of fish by families was 1.3%, 26.2% and 57%, respectively. Consumption of fish and n-3 (ω -3) poly-unsaturated fatty acids was associated with a lower risk of pancreatic cancer and other diseases (46-48). Our findings showed that most of the households used to eat tons, fizzy drinks and industrial juice rarely. In all households, buttermilk (60.8%) and milk, yoghurt or cheese (nearly 90%) was consumed daily. Wang H et al. determined greater intakes of total dairy foods; total low-fat/fat-free dairy foods, low-fat/skimmed milk and yoghurt were associated with smaller annualized increments in systolic blood pressure and a lower risk of projected hypertension incidence. Total dairy and total low-fat/fat-free dairy intakes were found to be inversely related to changes in diastolic blood pressure (49). In another study, the association between consumption of different types of dairy products and long-term changes in weight and waist circumference among American adults with an average follow-up of 13-years was examined (50). Our findings showed that 19.2% of the households consumed snacks weekly. The frequency of weekly egg, red meat, legumes and poultry consumption was 58.8%, 68.7%, and 49.6%, respectively; 80.6% of these was acceptable. Elevation of blood cholesterol concentrations has been recognized as a major risk factor for cardiovascular diseases and higher snack dietary pattern scores predicted a higher risk of hypercholesterolemia (51). Present study showed that, in both urban and rural areas more than 80% of the families consumed sugar daily. Most of the households consumed white bread and liquid oil for cooking.

In summary, although nutrition knowledge has been identified as an unreliable predictor of behavioral change, research suggests that nutrition knowledge is a critical factor in promoting favorable dietary and health changes. Results indicate that nutrition knowledge can impart favorable dietary and healthy behavioral changes.

CONCLUSION

A higher level of nutrition knowledge is associated with better practice and diet quality. Findings suggest that nutrition education should be integrated in household training programmes. Data from this study provided some useful information which could be used to develop areas of an intervention program for families.

Conflict of interests: none declared. Financial support: none declared Acknowledgements: The authors would like to thank all respected experts in the health departments of Urmia University of Medical Sciences and other professionals who have had sincere cooperation for conducting this study.



References

- Diet, nutrition and the prevention of chronic diseases: report of a joint WHO/ FAO expert consultation, 2003.
- Kamran A, Azadbakht L, Sharifirad G, Mahaki B, Sharghi A. Sodium intake, dietary knowledge, and illness perceptions of controlled and uncontrolled rural hypertensive patients. *Int J Hypertens* 2014;2014:245480.
- 3. Bano R AE, Fatima SB, Al-Shammari NA. A comparative study of Knowledge, Attitude, Practice of nutrition and non-nutrition student towards a balanced diet in Hail University. *IOSR Journal of Nursing and Health Science* 2013;2:29-36.
- Alaunyte I, Perry JL, Aubrey T. Nutritional knowledge and eating habits of professional rugby league players: does knowledge translate into practice? *J Int Soc Sports Nutr* 2015;12:18.
- Naeeni MM, Jafari S, Fouladgar M, et al. Nutritional Knowledge, Practice, and Dietary Habits among school Children and Adolescents. Int J Prev Med 2014;5:171-178.

- Bonaccio M, Di Castelnuovo A, Costanzo S, et al, Moli-sani Project I. Nutrition knowledge is associated with higher adherence to Mediterranean diet and lower prevalence of obesity. Results from the Moli-sani study. *Appetite* 2013;68:139-146.
- Golzarand M, Mirmiran P, Jessri M, et al. Dietary trends in the Middle East and North Africa: an ecological study (1961 to 2007). *Public Health Nutr* 2012;15:1835-1844.
- Noori MA, Ghiasvand R, Maghsoudi Z, et al. Evaluation of dietary pattern stability and physical activity in three consecutive generations of women. *Int J Public Health* 2015; 61:29-38.
- Zarei N, Ahmadi A: Nutrition Transition. An Intergenerational Comparison of Dietary Habits among Women of Shiraz. Iran J Public Health 2015;44:269-275.
- **10.** Larson N, Story M, Eisenberg ME, Neumark-Sztainer D. Secular Trends in Meal and Snack Patterns among Adolescents from 1999 to 2010.

J Acad Nutr Diet 2016;116:240-50.e2.

- Zarei M, Taib MN, Zarei F. Lifestyle factors and dietary intake of Iranian postgraduate students in Universiti Putra Malaysia (UPM). *Electron Physician* 2013;5:687-697.
- Aljefree N, Ahmed F. Association between dietary pattern and risk of cardiovascular disease among adults in the Middle East and North Africa region: a systematic review. *Food Nutr Res* 2015;59:274-86.
- Moran LJ, Grieger JA, Mishra GD, Teede HJ. The Association of a Mediterranean-Style Diet Pattern with Polycystic Ovary Syndrome Status in a Community Cohort Study. *Nutrients* 2015;7:8553-8564.
- 14. Afshin A MR, Khatibzadeh SH et al. The impact of dietary habits and metabolic risk factors on cardiovascular and diabetes mortality in countries of the Middle East and North Africa in 2010: a comparative risk assessment analysis. *BMJ Open* 2015;5:1-16.
- 15. Geaney F, Fitzgerald S, Harrington JM,

et al. Nutrition knowledge, diet quality and hypertension in a working population. *Preventive Medicine Reports* 2015, 2:105-113.

- **16.** Jessri M, Rashidkhani B: Dietary patterns and risk of gallbladder disease: a hospital-based case-control study in adult women. J Health Popul Nutr 2015;33:39-49.
- **17.** Mirmiran P, Azadbakht L, Azizi F. Dietary behaviour of Tehranian adolescents does not accord with their nutritional knowledge. *Public Health Nutr* 2007;10:897-901.
- Allafi AR, Alajmi F, Al-Haifi A. Survey of nutrition knowledge of physicians in Kuwait. *Public Health Nutr* 2013;16:1332-1336.
- **19. Cleland J**: A critique of KAP studies and some suggestions for their improvement. *Stud Fam Plann* 1973;4:42-47.
- 20. Chen ST, Soo KL, Azriani AR, et al. Nutrition knowledge, attitude and practice of teachers in rehabilitation centres in northern Malaysia. *Malays J Nutr* 2012;18:185-205.
- 21. Goh YM, Chua S: Knowledge, attitude and practices for design for safety. A study on civil & structural engineers. Accid Anal Prev 2016;93:260-6.
- 22. Kulkarni KK: KAP Studies Among Indian Antenatal Women. Can We Reduce the Incidence of Anemia? J Obstet Gynaecol India 2015;65:320-322.
- 23. Sharifirad G, Entezari MH, Kamran A, Azadbakht L. The effectiveness of nutritional education on the knowledge of diabetic patients using the health belief model. J Res Med Sci 2009;14:1-6.
- 24. Azizi F, Ghanbarian A, Momenan AA, et al. Prevention of non-communicable disease in a population in nutrition transition: Tehran Lipid and Glucose Study phase II. *Trials* 2009;10:5.
- 25. Kafeshani O, Sarrafzadegan N, Nouri F, Mohammadifard N. Major dietary patterns in Iranian adolescents: Isfahan Healthy Heart Program, Iran. *ARYA Atheroscler* 2015;11:61-68.
- 26. Abbastabar H, Roustazadeh A, Alizadeh A, et al. Relationships of Colorectal Cancer with Dietary Factors and Public Health Indicators: an Ecological Study. Asian Pacific Journal of Cancer Prevention 2015;16:3991-3995.
- 27. Safari A, Shariff ZM, Kandiah M, et al. Dietary patterns and risk of colorectal cancer in Tehran Province: a case-control study. *BMC Public Health* 2013;13:222.
- 28. Khosravi M, Sotoudeh G, Majdzadeh R, et al. Healthy and Unhealthy Dietary Patterns Are Related to Depression: A Case-Control Study. *Psychiatry Investig*

2015;12:434-442.

- **29.** Nouri Saeidlou S, Babaei F, Ayremlou P. Malnutrition, overweight, and obesity among urban and rural children in north of west Azerbijan, *Iran. J Obes* 2014;2014:541213.
- **30. Nouri Saeidlou S, Babaei F, Ayremlou P.** Children malnutrition in northwestern, central and southern regions of Iran: does geographic location matter? *Glob J Health Sci* 2014;6:36-41.
- Nouri Saeidlou S, Rezaiegoyjeloo F, Ayremlou P, Babaei F. Trend of Overweight and Obesity, Based on Population Study among School Children in North West of Iran: Implications for When to Intervene. MAEDICA 2015;10:214-220.
- **32.** Fallah F, Pourabbas A, Delpisheh A, Veisani Y, Shadnoush M. Effects of nutrition education on levels of nutritional awareness of pregnant women in Western iran. *Int J Endocrinol Metab* 2013;11:175-178.
- **33. Webb MC, Beckford SE.** Nutritional knowledge and attitudes of adolescent swimmers in trinidad and tobago. *J Nutr Metab* 2014;2014:506434.
- **34.** Noughani F, Bagheri M, Ramim T. Nutritional habits of mothers and children in the age group 0-4 years in Iran. *Ecol Food Nutr* 2014;53:410-418.
- **35. Keshteli AH, Feizi A, Esmaillzadeh A, et al**. Patterns of dietary behaviours identified by latent class analysis are associated with chronic uninvestigated dyspepsia. *Br J Nutr* 2015;113:803-812.
- 36. Sule SS, Onayade AA, Abiona TC, et al. Impact of nutritional education on nutritional status of under-five children in two rural communities of south-west Nigeria. Niger Postgrad Med J 2009;16:115-125.
- **37.** Azemati B, Heshmat R, Sanaei M, et al. Nutritional knowledge, attitude and practice of Iranian households and primary health care staff: NUTRIKAP Survey. J Diabetes Metab Disord 2013;12:12.
- **38. Hojhabrimanesh A, Akhlaghi M, Rahmani E, et al.** A Western dietary pattern is associated with higher blood pressure in Iranian adolescents. *Eur J Nutr* 2015:3.
- **39.** Farasati N, Siassi F, Koohdani F, et al. Western dietary pattern is related to premenstrual syndrome: a case-control study. *Br J Nutr* 2015: 114:2016-21.
- **40.** Naja F, Hwalla N, Itani L, et al. A Western dietary pattern is associated with overweight and obesity in a national sample of Lebanese adolescents (13-19 years): a cross-sectional study.

Br J Nutr 2015;114:1909-1919.

- **41.** Xu X, Hall J, Byles J, Shi Z. Dietary Pattern Is Associated with Obesity in Older People in China: Data from China Health and Nutrition Survey (CHNS). *Nutrients* 2015;7:8170-8188.
- 42. Shin S, Sung J, Joung H. A fruit, milk and whole grain dietary pattern is positively associated with bone mineral density in Korean healthy adults. *Eur J Clin Nutr* 2015;69:442-448.
- **43.** Sichert-Hellert W, Beghin L, De Henauw S, et al. Nutritional knowledge in European adolescents: results from the HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) study. *Public Health Nutr* 2011;14:2083-2091.
- 44. Sharma S, Dortch KS, Byrd-Williams C, et al. Nutrition-related knowledge, attitudes, and dietary behaviors among head start teachers in Texas: a crosssectional study. J Acad Nutr Diet 2013;113:558-562.
- **45.** Asghari G, Yuzbashian E, Mirmiran P, et al. Fast Food Intake Increases the Incidence of Metabolic Syndrome in Children and Adolescents: Tehran Lipid and Glucose Study. *PLoS One* 2015;10:e0139641.
- **46. Bryant J, Hanson M, Peebles C, et al.** Higher oily fish consumption in late pregnancy is associated with reduced aortic stiffness in the child at age 9 years. *Circ Res* 2015;116:1202-1205.
- **47. Del Brutto OH, Mera RM, Ha JE, et al.** Oily fish consumption is inversely correlated with cerebral microbleeds in community-dwelling older adults: results from the Atahualpa Project. *Aging Clin Exp Res* 2015;28:737-43.
- Hidaka A, Shimazu T, Sawada N, et al. Fish, n-3 PUFA consumption, and pancreatic cancer risk in Japanese: a large, population-based, prospective cohort study. Am J Clin Nutr 2015:102:1490-7
- **49. Wang H, Fox CS, Troy LM, et al.** Longitudinal association of dairy consumption with the changes in blood pressure and the risk of incident hypertension: the Framingham Heart Study. *Br J Nutr* 2015;114:1887-1899.
- 50. Wang H, Troy LM, Rogers GT, et al. Longitudinal association between dairy consumption and changes of body weight and waist circumference: the Framingham Heart Study. Int J Obes (Lond) 2014;38:299-305.
- **51.** Na L, Han T, Zhang W, et al A Snack Dietary Pattern Increases the Risk of Hypercholesterolemia in Northern Chinese Adults: A Prospective Cohort Study.*PLoS One* 2015;10:e0134294.