

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/266948461>

# Identification of medicinal plants affecting on headaches and migraines in Lorestan Province, West of Iran

Article in *Asian Pacific Journal of Tropical Medicine* · September 2014

DOI: 10.1016/S1995-7645(14)60261-3 · Source: PubMed

CITATIONS

32

READS

130

5 authors, including:



**Bahram Delfan**

Lorestan University of Medical Sciences

52 PUBLICATIONS 426 CITATIONS

[SEE PROFILE](#)



**Mahmoud Bahmani**

212 PUBLICATIONS 1,584 CITATIONS

[SEE PROFILE](#)



**Hassan Hassanzadazar**

Zanjan University of Medical Sciences

29 PUBLICATIONS 143 CITATIONS

[SEE PROFILE](#)



**Mahmoud Rafieian-kopaei**

Shahrekord University of Medical Sciences

314 PUBLICATIONS 4,015 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Aquapirins & choroid plexus [View project](#)



Evaluation of the effect of Green tea extracts on healing process in surgery and burning wounds [View project](#)



Contents lists available at ScienceDirect

## Asian Pacific Journal of Tropical Medicine

journal homepage: [www.elsevier.com/locate/apjtm](http://www.elsevier.com/locate/apjtm)

Document heading doi: 10.1016/S1995-7645(14)60261-3

## Identification of medicinal plants affecting on headaches and migraines in Lorestan Province, West of Iran

Bahram Delfan<sup>1</sup>, Mahmoud Bahmani<sup>1</sup>, Hassan Hassanzadazar<sup>2</sup>, Kourosh Saki<sup>3</sup>, Mahmoud Rafieian-Kopaei<sup>4\*</sup><sup>1</sup>Razi Herbal Medicines Research Center, Lorestan University of Medical Sciences, Khorramabad, Iran<sup>2</sup>Urmia University of Medical Sciences, Urmia, Iran<sup>3</sup>Shahid Beheshti University of Medical Sciences, Tehran, Iran<sup>4</sup>Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

## ARTICLE INFO

*Article history:*

Received 22 Jul 2014

Received in revised form 28 Aug 2014

Accepted 5 Sep 2014

Available online 29 Sep 2014

*Keywords:*

Pain

Headache

Medicinal plant

Lorestan province

Iran

## ABSTRACT

**Objective:** To identify medicinal plants which are used for headaches and migraines treatment in Lorestan province of Iran.**Methods:** Traditional medicinal herbs information was acquired from indigenous people with the cooperation of health centers of Doroud, Borujerd, Khorramabad, Aleshtar Poldokhtar, Aligoodarz, Nourabad and Kouhdasht cities. The prepared questionnaires were given to trained health volunteers. They attended in the villages and recorded people beliefs in herbal therapy by the questionnaires.**Results:** The results of this study showed that people used 15 herbs traditionally to treat headaches. Because of the importance of the medicinal plants in the study area, it is necessary to determine sociological studies the plenty of plant species.**Conclusions:** Because of the widespread use of traditional medicinal plants and high tendency to herbal medicine and traditional medicine, more extensive researches should be designed in several areas of pharmacy and pharmacology of medicinal plants to prepare proper information for pharmaceutical industries.

### 1. Introduction

Pain is the most common phenomena that force people to seek help from the health care system. Pain will affect various aspects of life, and not only caused stress and discomfort, but due to other stressors of individuals, such as treatment costs<sup>[1]</sup>.

In most epidemiological studies, headache symptoms are common with high frequency and different reasons such as tension or migraine. Headache is one of the ten most common causes for consultation with a doctor<sup>[2,3]</sup>.

Chronic tension-type headache is characterized by frequent attacks, often daily, non-pulsatile and bilateral pain behind the head without nausea, vomiting or visual disturbances. Pain is described as a tight band wrapped in head.

Migraine type headache is characterized by unilateral location, and pulsating quality with nausea, sensitivity to light and fatigue that will last for 7–4 h<sup>[4,5]</sup>. Migraine headache is the most common mental disorders with 12% prevalence in some studies<sup>[6,7]</sup>.

Stress, anxiety, assessment and response to small changes are important factors affecting the incidence of migraine<sup>[8,9]</sup>. Headache is one of the most prevalent complaints. Reports show 90% of people have a headache at least once a year<sup>[10,11]</sup>. Lorestan is one of the green provinces in the West of Iran which is located along the Zagros Mountains with unspoiled nature and rich plant

\*Corresponding author: Prof. Mahmoud Rafieian-Kopaei, Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran.

Tel: 0098 913 1811842

Fax: 0098 381 3349509

E-mail: [rafieian@yahoo.com](mailto:rafieian@yahoo.com)

Foundation Project: Supported by University of Medical Sciences of Lorestan province (Grant No. 2200/77).

flora. In this province, 151 native medicinal plants have been identified which used to treat a variety of diseases, traditionally<sup>[12]</sup>. This study was carried out to identify medicinal plants to treat headaches in this region.

## 2. Materials and methods

### 2.1. Studied region

Lorestan province is located in the west of Iran between 66° 51'–50° 3' east longitude from the Greenwich meridian and 32° 37'–34° 22' north latitude from the equator. This province has 4 different climates (semiarid, mild semi-humid, cold semi-humid and heights climate). The area of the Lorestan province is 28 300 ha.

Its minimum altitude is 330 m in the Zal Bridge and the maximum altitude is 4 050 m in the Oshtorankooch. This province has variable climate and this variety is completely obvious from north east to south west. Lorestan adjoins Hamedan and Markazi provinces in the north, and adjoins Isfahan in the east, Khuzestan in the south and Kermanshah and Ilam provinces in the west.

### 2.2. Methodology and plant collection

Traditional remedial information of this study

were obtained at 2007 to 2011, through interview and questionnaire and in cooperation with the Lorestan planning and management organization, Technology and Research Assistance of Lorestan Medical Sciences and benefiting indigenous information of people in provincial cities and also through cooperation of the treatment and health network in Doroud, Borujerd, Khoramabad, Aleshter, Pole Dokhtar, Aligudarz, Nourabad, and Kouhdasht cities in Lorestan province.

Questionnaires that had already been prepared by the directors of the health care system and were given to trained health volunteers. The questionnaire includes information about the location, characteristics of interviewer, native name and used parts of plant, mode of consumption, growing season and type of plants. Forms were completed by trained liaisons in the villages by asking old villagers. All information were recorded in tables.

## 3. Results

Fifteen medicinal plants described in ethnobotany study in Lorestan province have therapeutic effect against headache. Complete information of studied plants is listed in Table 1. Ethnobotanical information of these herbs included scientific name, family, local name, persian name, used parts of plant, using method and collection season.

**Table 1**

The ethnobotanical information of fifteen medicinal plants.

Scientific name	Family	Local name	Persian name	Used part of plant	Using method	Collection season	Application
<i>Echeveria elegans</i>	Crassulaceae	Pinome	Sagh	Seed	Herbal tea	Spring	Headache
<i>Alhagi persarum</i>	Fabaceae	Hashtarkhar	Khar shotor	Root	Herbal tea	Spring	Headache
<i>Allium haemanthoides</i>	Liliaceae	Ben sor	Piyaz yazdi	Leaves, tubers	Herbal tea	Spring, summer	Headache
<i>Althaea officinalis</i>	Malvaceae	Gole hiru	Khatmi	Flower, seed	Herbal tea	Spring, early summer	Headache
<i>Anchusa italica</i>	Boraginaceae	Gole gazo	Gaw Zaban	Flower, seed, root	Herbal tea	Spring, summer	Headache
<i>Artemisia annua</i>	Asteraceae	Khers dari	Darmaneh	Flower, stem	Herbal tea	All seasons	Headache
<i>Cichorium intybus</i>	Asteraceae	Cheghcheghe	Kasni	Root	Root decoction	All seasons	Headache
<i>Daphne mucronata</i>	Thymelaeaceae	Tolik	Mazarion	Bark, leaf	Roots incense leaves decoction	All seasons	Headache
<i>Falcaria vulgaris</i>	Apiaceae	Paghazou	Ghaziaghi	Leaf, flower, stem	The daily consumption of a cup of decoction	Spring, winter	Headache
<i>Ferula angulata</i>	Umbelliferae	Chavir	–	Flower, stem	Decoction	Spring, summer	Headache
<i>Matricaria recutita</i>	Compositae	Gole bayeneh	Babooneh	Petal	Herbal tea	Spring	Migraine
<i>Paliurus spina</i>	Rhamnaceae	Sipa	–	Leaf, stem	Decoction	Summer	Migraine
<i>Papaver rhoeasa</i>	Papaveraceae	Khashkhash	Khashkhash	Sap	Sap collection, drying and mixed with water and drink tea	Spring	Migraine
<i>Viola tricolor</i>	Umbelliferae	Gole banoushe	Banafsheh	Flower, branches	Herbal tea	Spring	Headache Migraine
<i>Ziziphus jujuba</i>	Rhamnaceae	Ano	Annab	Fruit, shell, core	Herbal tea	Autumn	Headache

#### 4. Discussion

For pain relief several chemical and synthetic drugs are supplied in pharmaceutical market with different side effects<sup>[13]</sup>. Side effects of medicinal plants are minimal. Comparison of this ethnobotanical study with other parts of Iran shows medicinal plants used for several purposes in Iranian traditional medicine.

Chicory (*Cichorium intybus*) has been named as a nerve tonic in Arasbaran. Chamomile (*Anthemis nobilis*) is used as an anti-inflammatory and anticonvulsant and oriental poppy (*Papaver orientale*) is used as an analgesic and hypnotic in this area of Iran<sup>[14]</sup>.

Camel thorn (*Alhagi*) is used for wound healing, kidney and bladder stones, hemorrhoids, jaundice in Sistan province and Jujube (*Zizyphus jujuba*) is used as laxative, respiratory diseases, kidney and bladder pain, blood purifier, for constipation in this area<sup>[15]</sup>.

Chamomile (*Anthemis austro-iranica*) is used to relieve cold natured and strengthens the heart in Kazeroon city. In this area Chicory (*Cichorium intybus* L.) is used as blood purifier to strengthen the stomach, and marshmallow (*Alcea aucheri* (Boiss.) Alef.) is used to prevent hair loss and laxatives, and camel thorn (*Alhagi*) is used for treatment of kidney stones<sup>[16]</sup>.

Armenian marshmallow (*Alcea aucheri* Boiss.) is used for sore throat in Cheshmeh Anjir region of Shiraz, and Alborzi poppy (*Papaver tennifolium* Boiss and Hohen ex Boiss.) is used as ecstasy and sedative<sup>[17]</sup>.

In Kerman, South marshmallow (*Alcea aucheri*) is used for treatment of colds and allergies, and used as laxative. Sagebrush (*Artemisia aucheri*) used as fever reliever, abdominal pain relievers and relieve weakness of body. Chicory plants used for removing heat stroke, liver failure and treatment of jaundice. Beef tongue (*Echium amoenum*) is used as a sedative and hypnotic. Jujube (*Zizyphus jujuba*) is used for cold treatment and is beneficial to improving the hot-tempered and used as laxative<sup>[18]</sup>.

Identification of various medicinal plants in different nations associated with study on their therapeutic and pharmaceutical effects have opened a new window for treatment of many diseases by researchers. Many species of medicinal plants are still unknown in the world used for treatment by indigenous people. Such information is orally transmitted between different ethnic groups from one generation to the next around the world, which can be gradually extinct in modern societies. First step to extract useful hidden information in such societies provides a detailed list of medicinal plants, uses and how to use these plants<sup>[19–33]</sup>.

Nowadays, use of natural and herbal medicines has

become increasingly common and expanding in most countries of the world which has led to extensive studies in the medical literature regarding the identification of medicinal plants. As a result, new resources have been introduced to produce useful drugs that have fewer side effects than synthetic drugs<sup>[26]</sup>.

Regarding the importance of the medicinal plants in the study area, it seems that sociological studies should investigate the frequency of plant species. Because of the widespread traditional use of medicinal plants and traditional medicine, much more extensive researches in various fields of pharmacy and pharmacology of medicinal plants are needed in this region. Researches will determine proper using methods of medicinal plants for personal and industrial policies through this valuable botanical information.

#### Conflict of interest statement

We declare that we have no conflict of interest.

#### Acknowledgements

This study was implemented in collaboration with the Management and Planning Organization and Research and Technology deputy of University of Medical Sciences of Lorestan province (Grant No. 2200/77). The authors give thanks to Marzieh Rashidipoor, Dr. Abbasi, Dr. Salarvand, Dr. Jamshidi, Dr. Maknali, Dr. Hejazi, Dr. Refahi, Dr. Roshani, Dr. Rezaei, Dr. Veisi, Dr. Mosaddegh, Vali Khorramabadi, Hassan Niknam, Salar Alizadeh, Bahman Hassanzadeh and all coworkers of Doroud, Borujerd, Khorramabad, Aleshtar, Poldokhtar, Aligoodarz, Nourabad and Kouhdasht Health Networks for cooperation in data collection. This study was also supported by Research deputy of Shahrekord University of Medical Sciences.

#### References

- [1] Shakeri R, Shaeiri MR, Roshan R. [The effect of biofeedback training on experimental pain according to perfectionism]. 2007; **11**(1): 43–59. Persian.
- [2] Rasmussen BK, Lipton RB. Epidemiology of headache. In: Olson J, Hansen PT, Welche KM, editors. *Headaches*. 2nd ed. Philadelphia: Lippincott Williams & Wilkins; 2000, p. 163–172.
- [3] Stovner Lj, Hagen K, Jensen R, Katsarava Z, Lipton R, Scher A, et al. The global burden of headache: a documentation of headache prevalence and disability worldwide. *Cephalalgia*

- 2007; **27**(3): 193–210.
- [4] Victor M, Roper AH, Victor M, Adams RD, Rooper AH. *Adams and Victor's principles of neurology*. 7th ed. New York: McGraw–Hill Professional; 2001, p. 175–192.
- [5] Natoli JL, Manack A, Dean B, Butler Q, Turkel CC, Stovner L, et al. Global prevalence of chronic migraine: a systematic review. *Cephalalgia* 2010; **30**(5): 599–609.
- [6] Stewart W, Linet M, Celentano DD, Van Natta M, Ziegler D. Age- and sex-specific incidence rates of migraine with and without visual aura. *Am J Epidemiol* 1991; **134**: 1111–1120.
- [7] Mathew NT. Pathophysiology, epidemiology, and impact of migraine. *Clin Cornerstone* 2001; **4**: 1–17.
- [8] Bener A, Swadi H, Qassimi E, Uduman S. Prevalence of headache and migraine in schoolchildren in the United Arab Emirates. *Ann Saudi Med* 1998; **18**(6): 522–524.
- [9] Lipton RB, Stewart WF, Diamond S, Diamond ML, Reed M. Prevalence and burden of migraine in the United States: data from the American Migraine Study II. *Headache* 2001; **41**: 646–657.
- [10] Modara F, Rostamkhani M. Prevalence of tension and migraine headaches among the students of Ilam Medical University. *J Ilam Univ Med Sci* 2008; **15**(4): 13–19.
- [11] Soltanzadeh A. *Neurologic disorders*. 1st ed. Iran: Mehrab; 1995.
- [12] Ahmadi S, Babakhanlu P, Karimifar MA. [Medicinal plants of Lorestan]. *Yafteh* 2010; **11**: 85–100. Persian.
- [13] Murray MD, Brater DC. Renal toxicity of the nonsteroidal anti-inflammatory drugs. *Annu Rev Pharmacol Toxicol* 1993; **33**: 435–465.
- [14] Zolfeghari E, Adeli I, Mozafarian V, Babaiy S, Habibi BG. [Identification of Arasbaran medicinal plants and ethnobotanical study of rural people knowledge]. *Iranian J Med Aromat Plants* 2012; **28**(3): 534–550. Persian.
- [15] Iranmanesh M, Najafi S, Yousefi M. Studies on ethnobotany of important medicinal plants in Sistan. *J Herbal Drugs* 2010; **1**: 58–65.
- [16] Dolatkahi M, Ghorbani-Nahoji M, Mehrafarin A, Amini Nejad GR, Dolatkahi A. Ethnobotanical study of medicinal plants in Kazeroon: identification, distribution and traditional uses. *J Med Plants* 2012; **11**: 163–178.
- [17] Sadeghi E, Borjian A. [Medicinal plants of Cheshme anger region (Shiraz)]. *J Res Plants Sci* 2013; **1**(7): 25, 42–59. Persian. [Online] Available from: [http://www.sid.ir/fa/VEWSSID/J\\_pdf/6000013912505.pdf](http://www.sid.ir/fa/VEWSSID/J_pdf/6000013912505.pdf) [Accessed on 25th April, 2014]
- [18] Sharififar F, Kouhpayeh A, Motaghi MM, Amirkhosravi A, Pourmohseni Nasab E, Khodashenas M. Study the ethnobotany of medicinal plants city in Sirjan, Kerman province, Iran. *J Herbal Drugs* 2010; **1**: 19–28.
- [19] Hazrat A, Nisar M, Shah J, Ahmad SH. Ethnobotanical study of some elite plants belonging to Dir, Kohistan valet, Khyber pukhtunkhwa, Pakistan. *Pak J Bot* 2011; **43**(2): 787–795.
- [20] Falsetto S. Medicinal properties of aromatic plant families. 2009. [Online] Available from: <https://suite.io/sharon-falsetto/1ays2e8c> [Accessed on 12nd March, 2014]
- [21] Bahmani M, Rafieian-Kopaei M, Avijgan M, Hosseini S, Golshahi H, Eftekhari Z, et al. Ethnobotanical studies of medicinal plants used by Kurdish owner's in south range of Ilam province, west of Iran. *American–Eurasian J Agric Environ Sci* 2012; **12**(9): 1128–1133.
- [22] Ghasemi Pirbalouti A, Momeni M, Bahmani M. Ethnobotanical study of medicinal plants used by Kurd tribe in Dehloran and Abdanan districts, Ilam Province, Iran. *Afr J Tradit Complement Altern Med* 2012; **10**(2): 368–385.
- [23] Bahmani M, Avijgan M, Hosseini SR, Bahmani E, Mehrzadi S. Traditional application of medicinal plants in southern area of Ilam province for treatment diseases and clinical syndromes in small ruminants. *J Herbal Drugs* 2011; **1**: 51–59.
- [24] Rafieian-Kopaei M. Medicinal plants and the human needs. *J HerbMed Pharmacol* 2012; **1**(1): 1–2.
- [25] Bahmani M, Eftekhari Z. An ethnoveterinary study of medicinal plants in treatment of diseases and syndromes of herd dog in southern regions of Ilam province, Iran. *Comp Clin Path* 2012; **22**: 403–407.
- [26] Ghasemi A. [Medicinal and aromatic plants]. Shahrekord: Islamic Azad University Press; 2011, p. 536. Persian.
- [27] Bahmani M, Farkhondeh T, Sadighara P. The anti-parasitic effects of *Nicotina tabacum* on leeches. *Comp Clin Pathol* 2012; **21**(3): 357–359.
- [28] Bahmani M, Karamati SA, Banihabib E, Saki K. Comparison of effect of nicotine and levamisole and ivermectin on mortality of leech. *Asian Pac J Trop Dis* 2014; **4**(Suppl 1): S477–S480.
- [29] Amirmohammadi M, Khajoenia S, Bahmani M, Rafieian-Kopaei M, Eftekhari Z, Qorbani M. *In vivo* evaluation of antiparasitic effects of *Artemisia abrotanum* and *Salvia officinalis* extracts on *Syphacia obvelata*, *Aspiculoris tetrapetra* and *Hymenolepis nana* parasites. *Asian Pac J Trop Dis* 2014; **4**(Suppl 1): S250–S254.
- [30] Eftekhari Z, Bahmani M, Mohsenzadegan A, Gholami–Ahangaran M, Abbasi J, Alighazi N. Evaluating the anti-leech (*Limnatis nilotica*) activity of methanolic extract of *Allium sativum* L. compared with levamisole and metronidazole. *Comp Clin Path* 2012; **21**: 1219–1222.
- [31] Bahmani M, Abbasi J, Mohsenzadegan A, Sadeghian S, Gholami–Ahangaran M. *Allium sativum* L.: the anti-immature leech (*Limnatis nilotica*) activity compared to Niclosomide. *Comp Clin Pathol* 2011; doi 10.1007/s00580–011–1380–7.
- [32] Bahmani M, Banihabib EKH, rafieian-Kopaei M. Comparison of disinfection activities of nicotine with copper sulphate in water containing *Limnatis nilotica*. *Kafkas Univ Vet Fak Derg* 2014; doi: 10.9775/kvfd.2014.11223.
- [33] Bahmani M, Saki K, Rafieian-Kopaei M, Karamati SA, Eftekhari Z, Jelodari M. The most common herbal medicines affecting *Sarcocystis* branches: a review study. *Asian Pac J Trop Biomed* 2014; **4**(12): 930–937.