

## Prevalence of Nocturnal Enuresis in School Aged Children The Role of Personal and Parents Related Socio-Economic and Educational Factors

Mahmoodzadeh, Hashem, MD; Amestejani, Morteza\*, MD; Karamyar, Mohammad, MD;  
Nikibakhsh, Ahmad-Ali, MD

Department of Pediatrics, Urmia University of Medical Sciences, Iran

Received: Jul 22, 2011; Accepted: Feb 17, 2012; First Online Available: Jan 02, 2013

### Abstract

**Objective:** Nocturnal enuresis is a common psychosocial concern for both parents and children. In the present study we have determined the prevalence of nocturnal enuresis in Urmia, Iran children and associated personal and familial factors with this problem.

**Methods:** A cross sectional epidemiological study for detection of nocturnal enuresis prevalence rate and evaluation of associated familial and personal factors in elementary school children (7-11 years old) from Urmia were investigated. The subjects were selected by cluster sampling method. Chi square test and logistic regression were used in univariate and multivariate respectively.

**Findings:** Of the 1600 questionnaires distributed, 918 (57%) were completed and included in the final analysis. The rest, which were not filled by parents and also those out of our study age range were excluded. Gender of the subjects was almost equally distributed (48.6% males and 51.4% females). Prevalence of nocturnal enuresis was 18.7% (n=172) and prevalence of day time incontinence was 5.5% (n=51). There was no significant gender difference between these two groups. Enuretics had crowded families, positive family history, low educational level of parents, jobless father, working mother, single parent, poor school performance, positive history of urinary tract infection (UTI).

**Conclusion:** Our results with enuresis prevalence and associated factors were comparable to other epidemiological studies from various countries. We found that Iranian families do not pay sufficient attention to their enuretic children.

*Iranian Journal of Pediatrics, Volume 23 (Number 1), February 2013, Pages: 59-64*

**Key Words:** Nocturnal enuresis; Diurnal enuresis; Prevalence; Children

### Introduction

Nocturnal enuresis can be defined as any intermittent incontinence while asleep in a child being at least five years old<sup>[1]</sup>. This disorder is a common, genetically complex and heterogeneous problem among children. This clinical problem is an important concern for family because it leads to considerable emotional distress and concern in

affected children, their parents and results in some psychological consequences, such as low self-esteem in children, some other psychological problems and specially low school success<sup>[2]</sup>. Bed wetting can be diagnosed at 5 year-olds and beyond; clinically, it is generally left untreated until the children are 7-8 years old<sup>[3]</sup>.

Most studies have been investigated to find the etiology of enuresis but most of them have found

\* Corresponding Author;

Address: School of Medicine, Urmia University, Urmia, Iran

E-mail: [mortezaamestejani@yahoo.com](mailto:mortezaamestejani@yahoo.com)

© 2013 by Pediatrics Center of Excellence, Children's Medical Center, Tehran University of Medical Sciences, All rights reserved.

only risk factors for it because this disorder has a multifactorial etiology [4-6].

We conducted a study to estimate the prevalence of nocturnal enuresis in Urmia, Iran. Different associated familial and personal factors with nocturnal enuresis were also studied.

## Subjects and Methods

Urmia is a mountainous city located in northwest Iran with a population of about 500000. A sample of 1600 students, 7-11 years old, representative for the primary school population of Urmia were included in the study. The city was classified into different regions. Every region was considered as a cluster. Seven clusters were selected randomly and two schools (a boys' and a girls' school) were selected in each cluster randomly. Finally fourteen schools were selected and questionnaires sent to the parents inviting them to come to school on special day after a week. 1142 (71%) of them came and we explained them our questionnaire; at the end, 974 of them filled the questionnaire completely under our supervision. Revising the questionnaires, we excluded children who were out of our age period range. At last 918 (57%) questionnaires were included in the study.

The items of the questionnaire were designed to collect four sections of information about children and their families (Fig. 1). The first section contained questions about the prevalence of enuresis and day time incontinence. Nocturnal enuresis was defined as any intermittent incontinence while asleep in a child being at least five years old, and day time incontinence any intermittent incontinence in day time in the same age range<sup>[1]</sup>. The second section required information on background data of the child such as age, gender, school performance, history of urinary tract infection (UTI), history of neonatal icterus, breast feeding lasting longer than one year,

history of febrile convulsion and birth weight. Educational performance was filled by parents due to elementary education office definition: good, moderate, and fail.

The third section was about data of the children's families such as history of enuresis in parents and siblings, parents' level of education and employment, divorced or dead parents, total number of family members, unplanned pregnancy and mother's smoking. Educational level of parents was divided into two groups: primary school or less as a low educational level and junior high school or more as a high educational level. Mother's smoking was defined as any smoking in pregnancy. Crowded family was defined as having 5 and more members. The fourth section dealt with treatment of enuresis such as visiting a physician and kind of treatment (medication, alarm treatment, fluid restriction and waiting for maturity). All analyses were performed with SPSS for Windows with  $P < 0.05$  considered to indicate significant association. Analyses were done by chi-square test for univariate and multivariate analysis, logistic regression to determine the significant predictive factors for nocturnal enuresis.

## Findings

Of the 1600 questionnaires distributed, 918 (57%) were included in analysis. These consisted of 453 boys and 465 girls. The overall prevalence of nocturnal and day time incontinence was 18.7% (n=172) and 5.5% (n=51) respectively (Table 1). The mean age was 8.6 in children with nocturnal enuresis and 9.09 in controls. There was a tendency for a steady decrease in the frequency of nocturnal enuresis with increasing age. There was no statistically significant difference in frequency of nocturnal enuresis in the two genders. Table 2 shows the frequency of nocturnal enuresis related to age and sex. Several familial factors that are related to nocturnal enuresis are shown in Table

**Table 1:** Prevalence of enuresis

Type of Enuresis	Boys n (%)	Girls n (%)	Total n (%)
Nocturnal Enuresis	95 (20.9)	77 (16.5)	172 (18.7)
Diurnal Enuresis	14 (3)	37 (7.9)	51 (5.5)

**Table 2:** The frequency of nocturnal enuresis in relation to age and gender

Age (years)	Boys n/N (%)*	Girls n/N (%)*	Total n/N (%)*
7	30/91 (32.9)	24/91 (26.3)	54 (31.3)
8	21/91 (23)	14/95 (14.7)	35 (20.3)
9	17/90 (18.8)	15/92 (16.3)	32 (18.6)
10	14/97 (14.4)	14/89 (15.7)	28 (16.2)
11	13/84 (15.4)	10/98 (10.2)	23 (13.3)
<b>Total</b>	95/453 (20.9)	77/465 (16.5)	--

n=number of enuretic children; N=total number in each age group; \*percentage in enuretic children

3. Crowded families, positive family history, low educational level of the parents, jobless father, working mother and single parent (divorced or widowed) were significantly higher in children with enuresis when compared to non-enuretics. Table 4 represents personal factors which were encountered more in enuretic group. Poor school performance, positive history of UTI, not breast feeding, and low birth weight were significantly higher in enuretics when compared to non-enuretics. Unplanned pregnancy, neonatal icterus, febrile convulsion, mother's smoking did not show any increase in enuretics when compared to non-enuretics. The percentage of children with enuresis attending a physician for treatment was 19.8%. Medication was the most (64.5%) preferred treatment for enuresis, waiting for maturity was the least (6.5%) preferred. Table 5 shows logistic regression test to determine the significant predictive factors for nocturnal enuresis.

## Discussion

This is a population-based study, which was aimed to detect prevalence and risk factors of nocturnal enuresis in Urmia, a city in northwest Iran. We obtained the prevalence of enuresis in 7-11 year-old children as 18.7%. In previous studies from other provinces, the prevalence of enuresis was reported 7.7%<sup>[7]</sup> and 6.8%<sup>[8]</sup> which is lower than ours, but in one province it was reported 17.5%<sup>[9]</sup> which is similar to our study. Also in children of a rural community in southwestern Nigeria and Turkey it has been reported 17.5%<sup>[10,2]</sup>. This is similar to our finding.

It is noticeable that the prevalence of nocturnal enuresis has been reported 8% for Taiwan children<sup>[11]</sup>, and 9.2% for Korean children<sup>[12]</sup>. This difference might be due to sociocultural variations between the countries and regions. Previous studies demonstrated that the prevalence of enuresis tended to decrease with increasing

**Table 3:** Family factors related to enuresis

Factors		Enuretics n (%)	Non-enuretics n (%)	P-value
<b>Educational level of mother</b>	Primary school or less	108 (62.8)	298 (39.9)	<0.001
	Junior high school or more	64 (37.2)	448 (60.1)	
<b>Educational level of father</b>	Primary school or less	83 (48.3)	225 (30.2)	<0.001
	Junior high school or more	89 (51.1)	521 (69.8)	
<b>Working Parents</b>	Mother	70 (40.6)	260 (34.8)	0.03
	Father	143 (83.1)	671 (89.9)	0.012
<b>Family history of enuresis</b>	Mother or father enuretic	53 (30.8)	68 (9.1)	<0.001
	Mother and father non- enuretic	119 (69.2)	678 (90.9)	
	Sibling enuretic	54 (31.4)	53 (7.1)	<0.001
	Sibling non-enuretic	118 (68.6)	693 (92.9)	
<b>Divorce</b>		22 (12.8)	56 (7.5)	0.03
<b>Parent death</b>		19 (11)	48 (6.4)	0.04
<b>Family number</b>	3	27 (15.7)	257 (34.5)	<0.001
	4	36 (20.9)	308 (41.3)	
	5	59 (34.3)	120 (16.1)	
	More than 5	50 (29.1)	61 (8.2)	
<b>Unwilling pregnancy</b>		17 (9.9)	69 (9.2)	NS
<b>Mother smoking</b>		13 (7.5)	50 (6.7)	NS

NS: Non-significant

**Table 4:** Personal factors that were related to enuresis

Factors		Enuretics	Non-enuretics	P Value
		N (%)	N (%)	
School performance	Good	62 (36)	575 (77.1)	<0.001
	Moderate	85 (49.4)	139 (18.6)	
	Fail	25 (14.5)	32 (4.3)	
History of UTI		36 (20.9)	47 (6.3)	<0.001
History of neonatal icter		62 (36)	243 (32.6)	NS
Not Breast feeding		48 (27.9)	128 (17.2)	0.001
History of febrile Convulsion		9 (5.2)	37 (4.6)	NS
Birth weight	< 2500gr	74 (43)	208 (27.9)	<0.001
	≥ 2500gr	98 (57)	538 (72.1)	

NS: Non-significant; UTI: Urinary tract infection

age<sup>[2,4]</sup>, similarly in the present study, 31.3% of the children were wetting their bed at age 7 whereas 13.3% of them were wetting their bed at age 11. As nocturnal enuresis is mostly expected to improve spontaneously, its decrease with age is thought to be mostly due to spontaneous improvement<sup>[4]</sup>.

In the present study, there was no statistically significant difference in prevalence of nocturnal enuresis between boys and girls (20.9% versus 16.5%). This result is also similar to most reports in the literature<sup>[13,14]</sup>. Although in some literature the prevalence of enuresis was more common in boys than in girls <sup>[2,5,10,11,13,15-20]</sup>. The present study showed that some familial factors were significantly associated with nocturnal enuresis.

Enuresis was more frequent among children with positive family history and this is consistent with the literature<sup>[2,4,5,11,20-22]</sup>. Some studies have reported parents of enuretics had low educational level<sup>[2,19,22]</sup>. We saw low educational level of fathers in 48.3% and 62.8% in mothers. Working mothers were found to have more enuretic

children than housewives. Jobless fathers were seen in 83.1% of enuretics. This was thought to be creating stress in their children. Similar results were seen in school age children in northwest Turkey and Istanbul<sup>[4,13]</sup>. Crowded family was another associated familial factor in enuretics. Also increased number of siblings was an effective factor in Turkish children<sup>[2]</sup> and children in Istanbul<sup>[13]</sup>. The present results also showed that single parent (divorced or parent death) is an important factor associated with enuresis. Our results correspond with those of the studies in Taiwan and Turkey<sup>[2,11]</sup>.

The personal factors encountered more in enuretics were: poor school performance, positive history of UTI, not breast feeding, and low birth weight. In our study poor school performance was comparable with other epidemiologic studies from various countries with poor school performance <sup>[2,4,21,23]</sup>. According to our findings, history of recurrent UTI was significantly associated with enuresis. 20.9% enuretics had positive history of UTI which is similar to that in Turkish and

**Table 5:** Multivariate analysis to determine the significant predictive factors for nocturnal enuresis

Factors	Coefficient Regression	P-value	Odd Ratio	95% Confidence interval
Parent s' history of enuresis	1.44	<0.001	3.14	1.93-5.10
Siblings' history of enuresis	1.9	<0.001	6.74	4.07-11.14
Not Breast feeding	1.21	<0.001	3.35	2.1-5.35
Positive UTI history	1.07	<0.001	2.92	1.6-4.16
Parent death	0.79	0.001	2.2	1.1-4.1
Not working father	0.63	0.02	1.87	1.08-3.25
Working mother	0.76	0.001	2.14	1.35-3.41
Low educational level of father	0.55	<0.001	3.14	1.93-5.1
Low educational level of mother	0.95	<0.001	2.6	1.72-3.93

European children<sup>[2,21]</sup>.

Beside other factors, the role of not breast feeding and low birth weight were found to be important in 27.9% and 43% of the enuretics compared with 17.2% and 27.9% non-enuretics. According to our findings unplanned pregnancy, neonatal icterus, febrile convulsion, or mother's smoking was not increased in enuretics when compared to non-enuretics. In the present study 19.8% (n=13) of the children were seen by a physician similar to that of 17.2% in Turkey<sup>[2]</sup>. In our study, the most commonly used methods for treatment were medication (64.5%). Also in Turkey<sup>[2]</sup> the most preferred treatment was medication (59.5%). In contrast, these results were not supported by studies from northwest Turkey<sup>[4]</sup>. Here only 19.8% of the cases used medication.

As a limitation to the study we did not consider the severity of enuresis and related factors. Furthermore, it is recommended that our findings should be confirmed by further studies with attention to severity factors related to enuresis.

## Conclusion

In summary, the prevalence of enuresis in Urmia was similar to studies in other countries having similar socioeconomic status. Our results represented enuresis associated with a lot of personal and familial factors which were comparable to other epidemiologic studies from various countries.

## Acknowledgment

This study has been approved and supported by Urmia Medical University.

**Conflict of Interest:** None

## References

1. Nevés T, von Gontard A, Hoebeke P, et al. The standardization of terminology of lower urinary tract function in children and adolescents: report

from the standardization committee of the international children's continence society. *J Urol* 2006;176(1):314-24.

2. Ozden C, L. Ozdal O, Altinova S, et al. Prevalence and Associated Factors of Enuresis in Turkish Children. *International Braz J Urol* 2007;33(2): 216-22.
3. C. Logan A, Lesperance F. Primary nocturnal enuresis: omega-3 fatty acids may be of therapeutic value. *Med Hypotheses* 2005;64(6): 1188-91.
4. Unalacak M, Söğüt A, Aktunç E, Demircan N, Altın R: enuresis nocturnal prevalence and risk factors among school age children in northwest turkey. *Eur J Gen Med* 2004;1(3):21-5.
5. De Sousa A, Kapoor H, Jagtap J, et al. Prevalence and factors affecting enuresis among primary school children. *Indian J Urol* 2007;23(4):354-7.
6. Howe AC, Walker CE. Behavioral management of toilet training, enuresis and encopresis. *Pediatr Clin North Am* 1992;39(3):413-32.
7. Azhir A, Frajzadegan Z, Adibi A, et al. An epidemiological study of enuresis among primary school children in Isfahan, Iran. *Saudi Med J* 2006; 27(10):1572-7.
8. Safarinejad MR: Prevalence of nocturnal enuresis, risk factors, associated familial factors and urinary pathology among school children in Iran. *J Pediatr Urol* 2007;3(6):443-52.
9. Ghahramani M, Basiry Moghadam M, Ghahramani A. Nocturnal Enuresis and its impact on Growth. *Iran J Pediatr* 2008;18(2):167-70.
10. Osungbade KO, Oshiname FO. Prevalence and perception of nocturnal enuresis in children of a rural community in southwest Nigeria. *Trop Doct* 2003; 3(4):234-6.
11. Chang P, Chen WJ, Tsai WY, et al. An epidemiological study of nocturnal enuresis in Taiwanese children. *BJU Int* 2001;87(7):678-81.
12. Lee SD, Sohn DW, Lee JZ, et al. An epidemiological study of enuresis in Korean children. *BJU Int* 2000; 85(7):869-73.
13. Gur E, Turhan P, Can G, et al. Enuresis: Prevalence, risk factors and urinary pathology among school children in Istanbul, Turkey. *Pediatr Int* 2004;46(1): 58-63.
14. Ozkan KU, Garipardic M, Toktamis A, et al. Enuresis prevalence and accompanying factors in schoolchildren: A questionnaire study from southeast Anatolia. *Urol Int* 2004;73(2):149-55.
15. Serel TA, Akhan G, Koyuncuoğlu HR, et al. Epidemiology of enuresis in Turkish Children. *Scand J Urol Nephrol*. 1997;31(6):537-9.
16. Oge O, Kocak I, Gemalmaz H. Enuresis: point prevalence and associated factors among Turkish children. *Turk J Pediatr* 2001;43(1):38-43.
17. Marugan de Miguelsanz JM, Lapena Lopez de Armentia S, et al. An epidemiological analysis of the sequence of bladder control and nocturnal Enuresis prevalence in the children of the province of leÓN. *An ESP Pediatr* 1996;44(6):561-7.

18. Butler RJ, Golding J, Northstone K. ALSPAC Study Team: Nocturnal enuresis at 7.5 years old: prevalence and analysis of clinical signs. *BJU Int* 2005;96(3):404-10
19. Cher TW, Lin GJ, Hsu KH. Prevalence of nocturnal enuresis and associated familial factors in primary school children in Taiwan. *J Urol* 2002;168(3): 1142-6.
20. Wei Wang Q, Guo Wen J, Kui Song D, et al. Bed-wetting in Chinese children: Epidemiology and predictive factors. *Neurourol Urodyn* 2007;26(4): 512-7.
21. Inan M, Tokuc B, Y. Aydiner C, et al. Personal characteristics of enuretic children: an epidemiological study from south-east Europe. *Urol Int* 2008;81(1):47-53.
22. Tai HL, Chang YJ, Chang SC, Chen GD, Chang CP, Chou MC: The epidemiology and factors associated with nocturnal enuresis and its severity in primary school children in Taiwan. *Acta Paediatr* 2007; 96(2):242-5.
23. Chang SSY, Ng CFN, Wong SN, Hong Kong Childhood Enuresis Study Group: behavioural problems in children and parenting stress associated with primary nocturnal enuresis in Hong Kong. *Acta Paediatr* 2002;91(4):475-9.

### Questionnaire

<b>Part 1</b>			
1-Does your child wet his/her bed during nighttime?	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
2-What is the frequency of bedwetting of your child?			
2 or more than 2 times per week <input type="checkbox"/>	1 time per week <input type="checkbox"/>	2 times per month <input type="checkbox"/>	1 time per month <input type="checkbox"/>
3-Did your child experience bedwetting during daytime?	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
<b>Part 2</b>			
4- Your child is	Male <input type="checkbox"/>	Female <input type="checkbox"/>	
5- Age of the child: age ____ years.			
6- How is the school performance of your child?	Good <input type="checkbox"/>	Moderate <input type="checkbox"/>	Fail <input type="checkbox"/>
7- History of recurrent urinary tract infection.	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
8- History of neonatal jaundice.	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
9- Did your child have breast feeding more than a year?	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
10- History of febrile convulsion.	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
<b>Part 3</b>			
11- History of enuresis in the parents.	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
12- History of enuresis in the siblings.	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
13- Education level of father.	Primary school or less <input type="checkbox"/>	Junior high school or more <input type="checkbox"/>	
14- Education level of mother.	Primary school or less <input type="checkbox"/>	Junior high school or more <input type="checkbox"/>	
15- Working father.	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
16- Working mother.	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
17- Dead father.	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
18- Dead mother.	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
19- Divorced parents.	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
20- Number of family members.	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>
21- Unwilling pregnancy.	Yes <input type="checkbox"/>	No <input type="checkbox"/>	more than 5 <input type="checkbox"/>
22- History of mother's smoking in pregnancy.	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
<b>Part 4</b>			
23- Has your child been treated for bedwetting?	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
24- What type of treatment did your child receive?			
Medications <input type="checkbox"/>	wait for maturity <input type="checkbox"/>	fluid restriction <input type="checkbox"/>	alarm treatment <input type="checkbox"/>

Fig. 1: Survey questionnaire