

A Systematic Review of the Most Important Medicinal Plants Native to Iran Effective on Testicular Morphology and Hormonal Testicular Function

Leila Zarei¹, Masoumeh Pourjabali² and Nasrollah Naghdi^{3*}, Siamak Naji-Haddadi², Ehsan Bahmani⁴

¹Department of Anatomical Sciences, Faculty of Medicine, Lorestan University of Medical Sciences, Khorramabad, Iran

²Department of Pathology, Faculty of Medicine, Urmia University of Medical Sciences, Urmia, Iran

³Biotechnology and Medicinal Plants Research Center, Ilam University of Medical Sciences, Ilam, Iran

⁴Leishmaniasis Research Center, Ilam University of Medical Sciences, Ilam, Iran

Abstract

Reproductive disorders and infertility are complicated disorders with different types of consequences including medical and psychological. Infertility affects both women and men. Evidence indicates that half of the infertility is due to male factor infertility. Using medicinal plants that are effective on the reproductive disorders and infertility is considered a suitable therapeutic approach. In this review, the search terms testicle, spermatogenesis, and medicinal plants were used to retrieve relevant publications in the databases including PubMed, Scopus, Google Scholar, and SID. This review focused on animal studies on the effect of medicinal plants on testicular morphology and hormonal testicular function. According to the findings, several herbal plants are effective on the testicular tissue, the physiology of the testicles, and hormonal testicular function. Bioactive compounds and antioxidant study has plants that have therapeutic effects on the tissue and testicular physiology.

Keywords: Testicle, spermatogenesis, testicular function, plant, Iran

INTRODUCTION

Humans are exposed to certain agents in living and work environments that can threaten their reproductive system function [1]. Reproductive disorders and infertility are complicated disorders with different types of consequences including medical and psychological [2]. Infertility affects both men and women. Evidence indicates that half of the infertility is due to male factor infertility [3]. Male reproductive system consists of testicles, genital tract, appendix glands, and external genitals. External genitals include endocrine and exocrine glands. Exocrine glands release sperm and endocrine glands do testosterone, i.e. male sex hormone [4]. Spermatogenesis refers to conversion of spermatogonia into spermatozoa in the testicles. In mature testicle, with development of sertoli cell cytoplasm around the germ cells, it is necessary to feed and maintain these cells and establish cellular communication throughout spermatogenesis process [5]. Impaired serum LH and FSH levels can lead to impaired serum testosterone level that causes certain changes in the spermatogenesis [6].

Urinary and reproductive system disorders is one of the problem of today's medical sciences which their treatment is important [7-10]. Medicinal plants are not only a food source [11,12] but are also considered an available source of drugs and supplements [13-18]. Historically, the research and use of medicinal plants and nature-based products has been on rise [9-11]. Because chemical drugs cause side effects, much research has been conducted to investigate the use of plant-based pharmaceutical products

[12-14]. Currently, development of various types of herbal drugs and their pharmaceutical effects are being studied [15,16]. Medicinal plants can exert potent antioxidant effects due to certain compounds such as polyphenols [15-18]. Because chemical drugs cause many side effects, the use of medicinal plants has attracted much attention in different countries including Iran where rich sources of plants are available [11-19]. The optimal effects of medicinal plants have been investigated in boosting fertility and treating different disorders such as prostate inflammation, oligospermia, low sperm motility, hormone imbalance, impotence, and varicocele [18,19]. In Iran, numerous medicinal plants are used to treat infertility and testicular disorders. The aim of this review was to report the findings of animal studies on medicinal plants' effects on hormonal testicular function and testicular tissue.

METHODOLOGY:

In this review, the search terms testicle, spermatogenesis, and medicinal plants were used to retrieve relevant publications in the databases including PubMed, Scopus, Google Scholar, and SID.

RESULTS:

The number of identified plants Based on the results, 24 medicinal plants are used as antimicrobials for treatment of testice. Additional information on medicinal plants Medicinal herbs with remedies effects on testice along with their km information are shown in Table 1.

Table 1. Herbal plants effective on physiology and histology of testis

Row	Botanical name	Family	Persian name	Main findings	Reference
1	<i>Centella asiatica</i> (L.) Urban.	Apiaceae	Ab-boshghabi	A study with male Wistar rats demonstrated that testosterone concentration was 16.4 ± 2.8 and 14.1 ± 0.09 nm/l in control and sham groups, respectively, and 15 ± 1.32 , 9.8 ± 0.05 , and 8.4 ± 0.31 nm/l in experimental groups treated with 10, 50, and 80 mg/ml <i>C. asiatica</i> extract, respectively.	[19]
2	<i>Ferula asafoetida</i>		Anghouzeh	An experimental study on rats demonstrated that testicular weight was not significantly different among different treatments. Increasing <i>F. asafoetida</i> dose caused decrease in the thickness of the seminiferous cell layers. The number of Leydig cells and Sertoli cells decreased in rats treated with 150 and 300 mg/ml <i>F. asafoetida</i> gum extract. <i>F. asafoetida</i> gum extract caused decrease in blood testosterone level compared to control treatment. This study showed that over 150 mg/ml <i>F. asafoetida</i> extract caused degradation of spermatogenesis tissue.	[20]
3	<i>Plantago major</i>	Liliaceae	Barhang	A study demonstrated that treatment with <i>P. major</i> leaf extract caused significant increase in testosterone concentration and sperm number compared to other group. <i>P. major</i> can help to boost infertility through improving spermatogenesis in diabetic mice.	[21]
4	<i>Dorema aucheri</i>		Bilhar	Four hundred mg/kg hydroalcoholic <i>D. aucheri</i> extract helped to improve spermatogenesis and increase sperm number in rats.	[22]
5	<i>Vitex Agnus Castus</i>	Verbenacea	Panj-angosht	An experimental study with mice demonstrated that <i>V. agnus castus</i> fruit extract caused decrease in LH and testosterone compared to control group. This observation was most marked when 365 mg/kg of the extract was administered.	[23]
6	<i>Allium cepa</i>	Liliaceae	Piaz	A study with adult male BALB/c mice showed that the mean internal diameter of seminiferous tubules in the experimental group increased significantly compared to the control group (56.32 ± 1.42 μ m vs. 31.27 ± 3.16 μ m). In addition to affecting the structure of the seminiferous tubules, treatment with raw <i>A. cepa</i> extract affected cell proliferation in the mice's seminiferous tubules and accelerated spermatogenesis compared to the controls.	[24]
7	<i>Dactylorhiza Lancibracteata</i>		Saalab	A study with adult male BALB/c mice showed that aqueous <i>D. lancibracteata</i> root extract caused increase in mean LH level in experimental group (0.6 ± 0.9) compared to placebo (0.39 ± 0.006) and control (0.38 ± 0.007) groups.	[25]
8	<i>Camellia sinensis</i>		Chaye-sabz	A study on adult male NMRI mice reported that hydroalcoholic <i>C. sinensis</i> extract caused significant decrease in sperm number, motility, viability, and normal morphology as well as significant decrease in seminiferous tubule diameter and germinal epithelial thickness in mice treated with sodium arsenite compared to control group.	[26]
9	<i>Launaea acanthodes</i>	Asteraceae	Charkheh	An experimental study with male Wistar rats showed that 300 mg/kg aqueous <i>L. acanthodes</i> caused significant increase in progressive motility percentage, normal morphology, and number of sperm in diabetic rats.	[27]
10			Khorma	A study with Wistar rats showed that serum testosterone level decreased significantly in groups treated with 0.05 and 0.1 g/kg extract compared to control group.	[28]
11	<i>Cinnamomum zeylanicum</i>		Darchin	An experimental study with mice reported that LH and FSH concentrations increased with increase in the dose of hydroalcoholic <i>C. zaylanicum</i> extract up to 200 and 400 mg/kg. Fifty and 100 mg/kg hydroalcoholic <i>C. zaylanicum</i> extract caused increase in serum testosterone level compared to control group.	[29]
12	<i>Berberis integerrima</i>	Berberidaceae	Zereshk-afshan	A study showed that 6-week treatment with 500 mg/kg body weight <i>B. integerrima</i> extract caused increase in testosterone level.	[30]
13	<i>Crocus sativus</i> L.		zafaran	A study with laboratory mice reported that primary spermatocytes number increased significantly in group treated with 100 mg/kg <i>C. sativus</i> for 48 hours.	[31]

Row	Botanical name	Family	Persian name	Main findings	Reference
14	<i>Zingiber officinale</i>		Zanjabil	A study on <i>Z. officinale</i> extract effect on male mice demonstrated that this extract caused significant increase in blood testosterone level, decrease in testicular malondialdehyde, and increase in antioxidant enzymes activities in the testicles ($P < 0.05$). According to the findings, testicular structure and spermatogenesis process were normal in the mice treated with <i>Z. officinale</i> extract.	[32]
15			Soya	Administration with 30, 60, and 120 mg/kg hydroalcoholic soybean extract caused no adverse effects on body weight, gonads structures, testicular function parameters, and histometric parameters. Soybean extract caused significant decrease in plasma LH level, decrease in germinal epithelium length and reproduction coefficient. Histological examinations showed no adverse effects on testicular tissue.	[33]
16	<i>Anethum g</i>		Shevid	An experimental study on hydroalcoholic <i>A. graveolens</i> seed extract effects on testicular tissue and spermatogenesis showed that testicular weight and volume decreased significantly in experimental groups compared to control group, but no significant difference was observed in the weight of epididym and defran cannal between the experimental groups and the control group. Seminiferous tubules diameters decreased significantly in the experimental groups compared to control group. The number of spermatogonia and primary spermatocytes decreased significantly compared to the control group. The number of sperm and spermatids decreased significantly compared to the control group only when 100 mg/kg <i>A. graveolens</i> extract was administered.	[34]
17	<i>Aloe vera</i>		Sabre-zard	An experimental study with rats demonstrated that 400 mg/kg body weight hydroalcoholic <i>A. vera</i> extract caused increase in the number of spermatogonia, Sertoli cells, and Leydig cells in the testis in the diabetic rats.	[35]
18	Calligonum		kaligonum	Thirty mg/kg calligonum extract caused significant improvement of sperm parameters such as motility, viability, and normal morphology in male mice.	[36]
19	<i>Cucurbita pepo</i> L.		Kado-tanbal	The use of 40% and 8% of <i>C. pepo</i> caused increase in the number of spermatogonia, primary spermatocytes, and spermatids.	[37]
20			Gerdou	A study with male Wistar rats demonstrated that 10-50 mg walnut kernel extract exerted effects on the progression and continuation of spermatogenesis till final steps and increase in Leydig cells and sperm in testicular tissue, but no remarkable effects on the seminiferous tubules tissue structure.	[38]
21	<i>Heracleum persicum</i>	Umbelliferae	Golpar	Alcoholic <i>H. persicum</i> extract (0.1, 0.2, and 0.4 g/kg) caused significant decrease in serum testosterone level in treated groups compared to control group.	[39]
22	<i>Carthamus tinctorius</i>		Golrang	Intraperitoneal administration with 1.4 and 8.2 mg <i>C. tinctorius</i> extract per day for 20 days caused significant increase in testosterone concentration.	[40]
23	<i>Capparis spinosa</i>	Capparidaceae	Logji	Thirty mg/kg hydroalcoholic <i>C. spinosa</i> extract caused significant increase in sperm motility and normal morphology in male rats.	[41]
24	<i>Salvia officinalis</i>	Labiatae	Maryam-goli	Treatment with 150 and 200 mg/kg <i>S. officinalis</i> extract caused significant increase in serum testosterone level, seminiferous tubules diameters, and sperm number in the tubes tunnels in male Wistar rats.	[42]

DISCUSSION

Infertility refers to the failure to conceive after 12 months or more of regular unprotected sexual intercourse. Global statistics show that 15% of young couples suffer from infertility. In Iran, the prevalence of infertility was reported to be 13-18.5%, which is approximately similar to the global statistics. The causes of infertility are widely various

including infectious causes such as genital tuberculosis, as well as inflammation, hormone causes, low sperm count, sterility, congenital causes, medications, trauma, and endocrine disorders. The mechanism by which these plants enhance the chance of fertility is not clear. However, these plants mostly have phenolic compounds especially flavonoid compounds which both have shown to possess

antimicrobial and anti-inflammatory activities [43-65]. Furthermore, high oxidative stress is also associated with increase in the chance of infertility and flavonoids have antioxidant activities [66-74]. The compounds with antioxidant activity are able to scavenge free radicals, preventing organ damage [75-87]. There are a lot of other plants which have flavonoid compounds with antioxidant activity [88-100]. Other natural methods are also accepted and acknowledged by people [101].

CONCLUSION:

Therefore, these plants may also have anti-fertility activity which worth examining. This is due to the active pharmaceutical ingredients in the herbal plants

AUTHORS CONTRIBUTION

This work was carried out in collaboration among all authors.

CONFLICT OF INTEREST

The authors declared no potential conflicts of interests with respect to the authorship and/or publication of this paper.

REFERENCES

- [1] Paul M, Frazier L. Reproductive Disorders. In: Levy BS, Wegman DH. Occupational Health: Recognizing and Preventing Work-Related Disease. 4th ed. Philadelphia: Lippincot Williams & Wilkins. 2000: 590-3.
- [2] Chandra, A. and Stephen, E.H. (1998). Impaired fecundity in the United States: 1982-1995. *Family Planning Perspectives*, 30(1): 34-42.
- [3] World Health Organization. (2000). WHO Manual for the Standardized Investigation and Diagnosis of the Infertile Couple. Cambridge: Cambridge University Press.
- [4] Ranjbar A, Zareini P. Human physiology: endocrinology & reproduction. 1st ed. Tehran: Ilia publication; 2007.
- [5] Rex AH, Luiz RF. Spermatogenesis and cycle of the seminiferous epithelium, In: Molecular Mechanisms in Spermatogenesis, C Yan Cheng (editor). New York; 2008; 1-15.
- [6] Zitzmann M. Effects of testosterone replacement and its pharmacogenetics on physical performance and metabolism. *Asian J Androl*. 2008; 10(3): 364- 72.
- [7] Tamadon MR, Saberi Far M, Soleimani A, Ghorbani R, Semnani V, Malek F, and Malek M. Evaluation of noninvasive tests for diagnosis of Helicobacter pylori infection in hemodialysis patients. *J Nephropathol*. 2013; 2(4): 249–253.
- [8] Tamadon MR, Jamshidi L, Soliemani A, Ghorbani R, Malek F, Malek M. Effect of different doses of folic acid on serum homocysteine level in patients on hemodialysis. *Iran J Kidney Dis*. 2011 Mar;5(2):93-6.
- [9] Asgari MR, Mohammadi E, FallahiKhoshknab M, Tamadon MR. Hemodialysis patients' perception from nurses' role in their adjustment with hemodialysis: A qualitative study. *koomesh*. 2011; 12 (4) :385-395.
- [10] Soliemani A, Nikouejad H, Tabatabaizade M, Mianehsaz E, Tamadon M. Effect of hydroxymethylglutaryl-CoA reductase inhibitors on low-density lipoprotein cholesterol, interleukin-6, and high-sensitivity C-reactive protein in end-stage renal disease. *Iran J Kidney Dis*. 2011 Jan;5(1):29-33.
- [11] Mahmoudi GA, Mahmoodnia L, Mirhosseini M. Medicinal plants with anti-poisoning toxicity of carbon tetrachloride: An overview of the most important medicinal plants native to Iran with anti-carbon tetrachloride toxicity. *J Global Pharma Technol* 2016; 8 (11): 17-20.
- [12] Mahmoudi GA, Mahmoodnia L, Mirhosseini M. A review on the most important medicinal herbs native to Iran with anti-acetaminophen toxicity. *J Global Pharma Technol* 2016; 8 (11): 12-16.
- [13] Amirmohammadi M, Khajoenia S, Bahmani M, Rafieian-Kopaei M, Eftekhari Z, Qorbani M. In vivo evaluation of antiparasitic effects of Artemisia abrotanum and Salvia officinalis extracts on Syphacia obvelata, Aspiculuris tetrapetra and Hymenolepis nana parasites. *Asian Pac J Trop Dis*. 2014; 4(Suppl 1): S250-S254.
- [14] Asadbeigi M, Mohammadi T, Rafieian-Kopaei M, Saki K, Bahmani M, Delfan B. Traditional effects of medicinal plants in the treatment of respiratory diseases and disorders: an ethnobotanical study in the Urmia. *Asian Pac J Trop Med*. 2014; 7(Suppl 1): S364-S368
- [15] Rezvaniarad A, Mardani M, Shirzad H, Ahmadzadeh SM, Asgari S, Naimi A, Mahmoudi GHA. Curcuma longa: A review of therapeutic effects in traditional and modern medical references. *J Chem Pharmac Sci* 2016; 9 (4): 3438-3448.
- [16] Mahmoudi, G.A., Almasi, V., Lorzadeh, N., Khansari, A. The reasons for using and not using alternative medicine in Khorramabad women, west of Iran. *J Pakistan Med Assoc* 2015; 65(6): 623-625.
- [17] Asadi SY, Parsaei P, Karimi M, Ezzati S, Zamiri A, Mohammadzadeh F, Rafieian-Kopaei M. Effect of green tea (Camellia sinensis) extract on healing process of surgical wounds in rat. *Int J Surg*. 2013;11(4):332-7.
- [18] Madihi Y., Merrikhi A., Baradaran A., Ghobadi S., Shahinfard N., Ansari R., Karimi A. Mesripour A., Rafieian-Kopaei M. Bioactive components and the effect of hydroalcoholic extract of Vaccinium myrtillus on postprandial atherosclerosis risk factors in rabbits. *Pak J Med Sci*. 2013; 29(1): 384-389.
- [19] Fakher Rahim, Ghasem Saki ; Majid Jasemi. The Effect of Centella Asiatica Alcoholic Extract on the Serum Levels of Testosterone FSH and LH in Male Wistar Rat. *Sabzevar Uni Med Sci J*. 2009; 16(1): 6-11
- [20] Ayoubi A, Arshami J, Valizadeh R, Mousavi z, Mousaei A. The effect of gum extract (Ferula assa-foetida) on blood parameters and histopathology of testes in male Wistar rats. *Iranian J Animal Sci Res*. 2013; 4(4): 310-315.
- [21] Nejati V, Khanshi F. Effect of hydroalcoholic extract of Plantago major Leaf on the Testis Morphology, Sperm Parameters and Testosterone Level in Streptozotocin-Induced Diabetic Rats. *Quarterly Horizon Med Sci*. 2014; 20(1): 49-55
- [22] Ahangarpour A1, Oroojan AA2, HeydariH. Effect of Hydro-Alcoholic Extract of Dorema Aucheri on Serum Levels of Testosterone, FSH and Sperm Count in Nicotinamide-STZ-Induced Diabetic Rat Models. *Zanjan Uni Med Sci J*. 2009; 21; 87; 22-31.
- [23] Nasri S, Oryian SH, Haeri-Rouhani A, Amin GHR, Taghi-Zadeh MA. Vitex fruit and its interaction with bromocriptine effects on LH and testosterone in male mice. *Hormozgan Uni Med Sci J*. 2005; 9(2): 113-118.
- [24] Nikravesh MR, Jalali M, Mohammadi SH. Effect of onion extract on testicular tissue of mice. *J Reprod Infertilit*. 2009; 10(4): 239-244.
- [25] Faraji Z1, Nikzad H*2, Parivar K3, Nikzad M. The effect of aqueous extract of Salep Tubers on the structure of testis and sexual hormones in male mice. *Jahrom Uni Med Sci J*. 2013; 11(1): 71-76.
- [26] Seyyed Mohammad Ali Shariatzadeh, MaryamMohammadi. Protective role of green tea (Camellia sinensis) hydroalcoholic extract on sperm parameters and testicular tissue in NMRI mice exposed to sodium arsenite. *Birjand Uni Med Sci J*. 2014; 432-443.
- [27] Rahbarian R., Sepehri Moghadam H., Sadoughi SD. Effect of Aqueous Extract of Launaea acanthodes on Testicular Tissue and Sperm Parameters in Alloxan-Induced Diabetic Rats. *Quarterly Horizon Med Sci*. 2015; 21(1): 21-29.
- [28] Mokhtari M Sharifi E and Moghadamnia D. Effects of alcoholic extract of date on changes in testicular tissue and the amount of FSH, LH and testosterone in male rats. *J Basic Med Sci*. 2006; 9(4): 265-271.
- [29] Modaresi M, Messripour M, Toghiani M, Rajaii RA. Effect of hydroalcoholic extract of Cinnamon zeylanicum (Bark) on mice pituitary-testis axis. *J Gorgan Uni Med Sci*. 2010; 12 (1):15-19.
- [30] Ashraf H, Khaneshi F, Rafiee Raki F, Nejati V. Evaluation of Aqueous Extract of Berberis Integerrima Root on the Testis Tissue and Testosterone Levels in Streptozotocine (STZ) Induced Diabetic Rats. *Qom Univ Med Sci J*. 2013; 7 (4):28-35.

- [31] M. Modaresi, M. Messripor, M. Asadi Morghmaleki and M.K. Hamadani. The effect of Saffron extract on testis tissue. *Iranian J Med Aromatic Plants*. 2008; 24(2): 237-243.
- [32] Amoughli Tabrizi, B., Khakpour, M. Protective Effects of Ginger (*Zingiber officinale*) Rhizome Extract on Heat-Induced Testicular Damage in the Mouse. *J Vet Clin Pathol*. 2013; 27(7(3): 183-192.
- [33] Ahsannia N, Zahiri S, Mahjour A, Tavasoli A, Safarpour D. Histometric and Histopathologic Study of Soybean Hydroalcoholic Extract on Testicular Tissue and Pituitary-Gonad Axis in Adult Male Rat. *J Mazandaran Univ Med Sci*. 2011; 21(85): 92-98.
- [34] Salamat-Manesh M, Shiravi AH, Heydari-Nasrabadi M. The effect of alcoholic extract of fennel seeds (*Anethum graveolens*) on the testes and spermatogenesis in rats Male Wistar. *J Biol*. 2009; 1(2): 23-30.
- [35] Farhangdoost F, Jafari Barmak M, Hemayatkhah Jahromi V, Azizi A, Mahmoodi R, Keshavarzi E, Naraki M. Aloe Vera Extract Effect on Sperm Quality and Testicular Tissue of Rats Induced by Cadmium Chloride. *Yasuj Uni Med Sci J*. 2014; 84: 19(1): 47-55.
- [36] Asgari-Jahromi M, Movahedin M, Amanlou M, Moli SJ, Mazaheri Z, Batouli H. The effect of Calligonum extract the sperm and testis cell death in older planning. *Biol Pathol*. 2013; 16(1): 41-54.
- [37] Mohsen pour M, Shariati M, Esfandiari A. Effect of Cucurbita pepo L. Fruit Pulp on Testis Histological Changes in Adult Rats. *Yasuj Uni Med Sci J*. 2014; 89: 19(6): 518-528.
- [38] Abedinzade M, Mokhtari M, Zarbakhsh S, Nasseran SN. The Effect of the Alcoholic Extract of Walnut on the Testis Tissue of Adult Male Rats. *Yasuj Uni Med Sci J*. 2012; 69: 17(3): 215-224.
- [39] Fatemeh Barzegari Firouzabadi and Mahboubeh Mirhosseini. Effect of Persian hogweed (*Heracleum persicum*) on the morphological changes in mice testes and the level of hormone testosterone. *Razi j Med Sci*. 2012; 19(9): 18-24.
- [40] Modaresi M. Effect of alcoholic extract of safflower (*Carthamus tinctorius*) on the pituitary-gonadal tissue in the testes of mice. *Zanjan Uni Med Sci J*. 2005; 13(53): 1-7.
- [41] Mohammadi J, Chatroz B, Delaviz H. The Effect of Hydroalcoholic Extract of Capparis Spinosa on Quality of Sperm and Rate of Testosterone Following Induction of Diabetes in Rats. *J Isfahan Med Sch*. 2014; 31(264): 2042-52.
- [42] Ahmadi R, Balali S, Tavakoli P, Mafi M, Haji G R. The effect of hydroalcoholic leaf extract of *Salvia officinalis* on serum levels of FSH, LH, testosterone and testicular tissue in rats. *Feyz*. 2013; 17(3): 225-231.
- [43] Asgari S, Rafieian-Kopaei M, Shamsi F, Najafi S, Sahebkar A. Biochemical and histopathological study of the anti-hyperglycemic and anti-hyperlipidemic effects of cornelian cherry (*Cornus mas L.*) in alloxan-induced diabetic rats. *J Complement Integr Med*. 2014; 11(2):63-9.
- [44] Bahmani M, Sarrafchi A, Shirzad H, Rafieian-Kopaei M. Autism: Pathophysiology and promising herbal remedies. *Curr Pharm Des*. 2016; 22(3):277-285.
- [45] Bahmani M, Shirzad H, Mirhosseini M, Mesripour A, Rafieian-Kopaei M. A Review on Ethnobotanical and Therapeutic Uses of Fenugreek (*Trigonella foenum-graceum L.*). *J Evid Based Complementary Altern Med*. 2015 Apr 27. pii: 2156587215583405. [Epub ahead of print].
- [46] Karamati SA, Hassanzadazar H, Bahmani M, Rafieian-Kopaei M. Herbal and chemical drugs effective on malaria. *Asian Pac J Trop Dis*. 2014; 4(Suppl 2): 599-601.
- [47] Khosravi-Boroujeni H, Mohammadifard N, Sarrafzadegan N, Sajjadi F, Maghroun M, Khosravi A, Alikhasi H, Rafieian M, Azadbakht L. Potato consumption and cardiovascular disease risk factors among Iranian population. *Int J Food Sci Nutr*. 2012; 63(8):913-20.
- [48] Khosravi-Boroujeni H, Sarrafzadegan N, Mohammadifard N, Sajjadi F, Maghroun M, Asgari S, Rafieian-Kopaei M, Azadbakht L. White rice consumption and CVD risk factors among Iranian population. *J Health Popul Nutr*. 2013; 31(2): 252-61.
- [49] Madihi Y, Merrikhi A, Baradaran A, Rafieian-kopaei M, Shahinfard N, Ansari R, Shirzad H, Mesripour A. Impact of sumac on postprandial high-fat oxidative stress. *Pak J Med Sci*. 2013; 29 (1): 340-345.
- [50] Ghaed F, Rafieian-Kopaei M, Nematbakhsh M, Baradaran A, Nasri H. Ameliorative effects of metformin on renal histologic and biochemical alterations of gentamicin-induced renal toxicity in Wistar rats. *J Res Med Sci*. 2012; 17 (7): 621-625.
- [51] Gharipour M, Ramezani MA, Sadeghi M, Khosravi A, Masjedi M, Khosravi-Boroujeni H. et al. Sex based levels of C reactive protein and white blood cell count in subjects with metabolic syndrome: Isfahan Healthy Heart Program. *J Res Med Sci*. 2013; 18: 467-72.
- [52] Heidarian E, Rafieian-Kopaei M. Protective effect of artichoke (*Cynara scolymus*) leaf extract against lead toxicity in rat. *Pharm Biol*. 2013; 51(9):1104-9.
- [53] Hosseini-asl K., Rafieian-kopaei M. Can patients with active duodenal ulcer fast Ramadan? *Am J Gastroenterol*. 2002; 97(9): 2471-2472.
- [54] Kafash-Farkhad N, Asadi-Samani M, Rafieian-Kopaei M. A review on phytochemistry and pharmacological effects of Prangos ferulacea (L.) Lindl. *Life Sci J*. 2013; 10(8s):360-367
- [55] Bahmani M, Shirzad H, Rafieian S, Rafieian-Kopaei M. Silybum marianum: Beyond Hepatoprotection. *J Evidence-Based Complement Alternative Med*. 2015; 20(4): 292-301.
- [56] Bahmani, M., Rafieian-Kopaei, M. Medicinal plants and secondary metabolites for leech control. *Asian Pac J Trop Dis*. 2014; 4(4): 315-316.
- [57] Delfan B, Bahmani M, Rafieian-Kopaei M, Delfan M, Saki K. A review study on ethnobotanical study of medicinal plants used in relief of toothache in Lorestan Province, Iran. *Asian Pac J Trop Dis*. 2014; 4(Suppl 2): 879-884.
- [58] Ebrahimie M, Bahmani M, Shirzad H, Rafieian-Kopaei M, Saki K. A Review Study on the Effect of Iranian Herbal Medicines on Opioid Withdrawal Syndrome. *J Evid Based Complementary Altern Med*. 2015; 20(4):302-9.
- [59] Ganji-Arjenaki M, Rafieian-Kopaei M. Probiotics are a good choice in remission of inflammatory bowel diseases: A Meta Analysis and systematic review. *J Cellular Physiol*. 2017 15. PubMed PMID: 28294322.
- [60] Murray FT, Cameron DF, Orth JM, Katovich MJ. Gonadal dysfunction in the spontaneously diabetic BB rats: alteration of the testes morphology, serum testosterone and LH. *Horm Metab Res*. 1985; 17(10):495-501.
- [61] Vignon F, Le Faou A, Montagnon D, Pradignac A, Cranz C, Winiszewsky P, et al. Comparative study of semen in diabetic and healthy men. *Diabete Metab*. 1991; 17(3):350-4.
- [62] Shayganni E, Bahmani M, Asgari S, Rafieian-Kopaei M. Inflammaging and cardiovascular disease: management by medicinal plants. *Phytomedicine*. 2016; 23: 1119-1126.
- [63] Bahmani M, Banihabib E Rafieian-Kopaei M, Gholami-Ahangaran M. Comparison of Disinfection Activities of Nicotine with Copper Sulphate in water Containing *Limnatis nilotica*. *Kafkas Univ Vet Fak Derg*. 2015; 21 (1): 9-11
- [64] Amidi N, Moradkhani S, Sedaghat M, Khiripour N, Larki-Harchegani A, Zadkhosh N, et al. Effect of green tea on inflammation and oxidative stress in cisplatin-induced experimental liver function. *J HerbMed Pharmacol*. 2016; 5(3):99-102.
- [65] Bahmani, M., Sarrafchi, A., Shirzad, H., Rafieian-Kopaei, M. Autism: Pathophysiology and promising herbal remedies. *Current Pharmaceutical Design*. 2016; 22(3): 277-285.
- [66] Baradaran A, Nasri H, Nematbakhsh M, Rafieian-Kopaei M. Antioxidant activity and preventive effect of aqueous leaf extract of Aloe Vera on gentamicin-induced nephrotoxicity in male Wistar rats. *Clinica Terapeutica*. 2014; 165(1):7-11.
- [67] Baradaran A, Nasri H, Rafieian-Kopaei M. Erythropoietin and renal protection. *Daru*. 2013; 20; 21(1):78.
- [68] Nasri H, Rafieian-Kopaei M. Medicinal plants and antioxidants: Why they are not always beneficial? *Iranian J Public Health*. 2014. 43(2):255-257.
- [69] Nasri H, Rafieian-Kopaei M. Medicinal plants and new concerns in statin consumption. *Iranian J Pub Health*. 2013; 42(9): 1071-1072.
- [70] Nasri H, Rafieian-Kopaei M. Oxidative stress and aging prevention. *Int J Prev Med*. 2013; 4(9): 1101-1102.
- [71] Baradaran A, Nasri H, Rafieian-Kopaei M. Oxidative stress and hypertension: Possibility of hypertension therapy with antioxidants. *J Res Med Sci*. 2014; 19(4):358-67.

- [72] Nasri H, Rafieian-Kopaei M. Protective effects of herbal antioxidants on diabetic kidney disease. *J Res Med Sci*. 2014;19(1):82-3.
- [73] Nasri H, Rafieian-Kopaei M. Tubular kidney protection by antioxidants. *Iran J Public Health*. 2013; 42(10):1194-1196.
- [74] Taghikhani A, Afrough H, Ansari-Samani R, Shahinfard N, Rafieian-Kopaei M. Assessing the toxic effects of hydroalcoholic extract of *Stachys lavandulifolia* Vahl on rat's liver. *Bratisl Lek Listy*. 2014;115(3):121-4.
- [75] Rafieian-Kopaei M, Baradaran A. Plants antioxidants: From laboratory to clinic. *J Nephropathol*. 2013; 2(2): 152-153.
- [76] Rafieian-Kopaei M, Shahinfard N, Rouhi-Boroujeni H, Gharipour M, Darvishzadeh-Boroujeni P. Effects of *Ferulago angulata* extract on serum lipids and lipid peroxidation. *Evidence-Based Complementary and Alternative Medicine*; 2014 (2014), Article ID 680856, 4 pages <http://dx.doi.org/10.1155/2014/680856>.
- [77] Nasri H, Shabnam Hajian Sh, Ahmadi A, Baradaran A, Kohi G, Nasri P, Rafieian-Kopaei M. Ameliorative Effect of Green Tea Against Contrast-induced Renal Tubular Cell Injury, *Iran J KID DIS*. 2015;9:421-6.
- [78] Nasri H, Shirzad H, Baradaran A, Rafieian-kopaei M. Antioxidant plants and diabetes mellitus. *J Res Med Sci*. 2015; 20:491-50
- [79] Rafieian-Kopaei M, Nasri H. Comment on: Is the renoprotective effect of erythropoietin in chronic kidney disease a myth? *J Formos Med Assoc*. 2014;113(1):62.
- [80] Rafieian-Kopaei M, Baradaran A, Rafieian M. Oxidative stress and the paradoxical effects of antioxidants. *J Res Med Sci*. 2013; 18(7): 628.
- [81] Sarrafchi A, Bahmani M, Shirzad H, Rafieian-Kopaei M. Oxidative stress and Parkinson's disease: New hopes in treatment with herbal antioxidants. *Curr Pharm Des*. 2016; 22(2): 238 – 246.
- [82] Saki K, Bahmani M, Rafieian-Kopaei M, Hassanzadazar H, Dehghan K, Bahmani F, Asadzadeh J. The most common native medicinal plants used for psychiatric and neurological disorders in Urmia city, northwest of Iran. *Asian Pac J Trop Dis*. 2014; 4(Suppl 2): 895-901.
- [83] Saki K, Bahmani M, Rafieian-Kopaei M. The effect of most important medicinal plants on two important psychiatric disorders (anxiety and depression)-a review. *Asian Pac J Trop Med*. 2014; 7(Suppl 1): 34-42.
- [84] Delfan B, Bahmani M, Hassanzadazar H, Saki K, Rafieian-Kopaei M. Identification of medicinal plants affecting on headaches and migraines in Lorestan Province, West of Iran. *Asian Pac J Trop Med*. 2014; 7(Suppl 1): 376-379.
- [85] Tavakolli N, Ghanadian M, Asghari Gh, Sadraei H, Asgari-Borjloul N., Tabakhian M. Development of a validated HPLC method for determination of an active component in *Pycnocycla spinosa* and tablets prepared from its extract. *J Herbmed Pharmacol*. 2017;6(1):37-42.
- [86] Rabie Z, Gholami M, Rafieian-Kopaei M. Antidepressant effects of *Mentha pulegium* in mice. *Bangladesh J Pharmacol*. 2016; 11(3): 711-715 doi:<http://dx.doi.org/10.3329/bjp.v11i3.27318>.
- [87] Rouhi-Boroujeni H, Heidarian E, Rouhi-Boroujeni H, Deris F, Rafieian-Kopaei M. Medicinal Plants with multiple effects on cardiovascular diseases: a systematic review. *Curr Pharm Des*. 2017; 23(7): 999 – 1015. DOI: 10.2174/1381612822666161021160524
- [88] Sharafati-Chaleshtori R, Shirzad H, Rafieian-Kopaei M, Soltani A. Melatonin and human mitochondrial diseases. *J Res Med Sci* 2016;21:138.
- [89] Gupta A, Shaikh AC, Chaphalkar SR. Aqueous extract of *Calamus rotang* as a novel immunoadjuvant enhances both humoral and cell mediated immune response. *J Herbmed Pharmacol*. 2017;6(1):43-48.
- [90] Shirzad H, Shahrani M, Rafieian-Kopaei M. Comparison of morphine and tramadol effects on phagocytic activity of mice peritoneal phagocytes in vivo. *Int Immunopharmacol*. 2009; 9 (7-8):968-70.
- [91] Kooti W, Ghasemiboroon M, Ahangarpour A, Hardani A, Amirzargar A, Asadi-Samani M, et al. The effect of hydro-alcoholic extract of celery on male rats in fertility control and sex ratio of rat offspring. *Journal of Babol University of Medical Sciences*. 2014;16(4):43-9.
- [92] Jivad N, Bahmani M, Asadi-Samani M. A review of the most important medicinal plants effective on wound healing on ethnobotany evidence of Iran. *Der Pharmacia Lettre*. 2016;8(2):353-7.
- [93] Parsaei P, Bahmani M, Karimi M, Naghdi N, Asadi-Samani M, Rafieian-Kopaei M. A review of analgesic medicinal plants in Iran. *Der Pharmacia Lettre*. 2016;8(2):43-51.
- [94] Kooti W, Ghasemiboroon M, Asadi-Samani M, Ahangarpour A, Abadi MNA, Afrisham R, et al. The effects of hydro-alcoholic extract of celery on lipid profile of rats fed a high fat diet. *Advances in Environmental Biology*. 2014;8(4):325-30.
- [95] Jivad N, Asadi-Samani M, Moradi MT. The most important medicinal plants effective on migraine: A review of ethnobotanical studies in Iran. *Der Pharma Chemica*. 2016;8(2):462-6.
- [96] Shirzad H, Taji F, Rafieian-Kopaei M. Correlation between antioxidant activity of garlic extracts and WEHI-164 fibrosarcoma tumor growth in BALB/c mice. *J Med Food*. 2011; 14(9):969-74.
- [97] Shaygani E, Bahmani M., Asgary S, Rafieian-Kopaei M. Inflammaging and cardiovascular disease: Management by medicinal plants, *Phytomedicine*. 2015, <http://dx.doi.org/10.1016/j.phymed.2015.11.004>
- [98] Rabie Z, Gholami M, Rafieian-Kopaei M. Antidepressant effects of *Mentha pulegium* in mice. *Bangladesh J Pharmacol*. 2016; 11(3): 711-715 doi:<http://dx.doi.org/10.3329/bjp.v11i3.27318>.
- [99] Sharafati-Chaleshtori R, Shirzad H, Rafieian-Kopaei M, Soltani A. Melatonin and human mitochondrial diseases. *J Res Med Sci* 2016;21:138.
- [100] Rouhi-Boroujeni H, Heidarian E, Rouhi-Boroujeni H, Deris F, Rafieian-Kopaei M. Medicinal Plants with multiple effects on cardiovascular diseases: a systematic review. *Curr Pharm Des*. 2017; 23(7): 999 – 1015. DOI: 10.2174/1381612822666161021160524
- [101] Ganji-Arjenaki M, Rafieian-Kopaei M. Probiotics are a good choice in remission of inflammatory bowel diseases: A Meta analysis and systematic review. *Journal of Cellular Physiology*. 2017 Mar 15. PubMed PMID: 28294322.