


Contents lists available at [ScienceDirect](#)

International Journal of Gerontology

journal homepage: www.ijge-online.com

Original Article

A Theoretically Based Behavioral Nutrition Intervention for Elderly Women: A Cluster Randomized Controlled Trial

Jamileh Amirzadeh Iranagh^{a*}, Seyedeh Ameneh Motalebi^b, Fatemeh Mohammadi^b^a Faculty of Health, Urmia University of Medical Sciences, Urmia, Iran, ^b Faculty of Nursing and Midwifery, Qazvin University of Medical Sciences, Qazvin, Iran

ARTICLE INFO

Article history:

Received 25 April 2017

Received in revised form

1 August 2017

Accepted 14 August 2017

Available online xxx

Keywords:

nutritional intervention,
elderly woman,
Health Belief Model

SUMMARY

Background: Nutritional behavior is one of the main criteria for determining health in the elderly. Health Belief Model (HBM) is most commonly used theory in the interventional programs. This study was aimed to determine efficacy of HBM-based nutritional educational intervention on knowledge, belief and behavior of the elderly women.

Methods: The sample of this randomized controlled trial study was 100 elderly women who attended to four health care centers of Urmia, Iran. Data were collected using self-administered questionnaire included demographic characteristics and some HBM constructs. The experiment group participated to HBM-based trainings classes twice per week for 12 weeks. The educational strategies were consisted of lecture, focus-group discussion, and video showing. The questionnaire was completed before and after 3 and 6 months intervention. Data were analyzed using two way repeated measure ANOVA at the significant level of $\alpha < 0.05$.

Results: The mean scores of benefit, severity, susceptibility, and self-efficacy were not different between the two groups before the educational training ($p > 0.05$). However, the aforementioned scores changed significantly between two group after 3 months intervention and follow-up ($p < 0.001$). Furthermore, intervention group improved significantly on HBM constructs through three times measurements ($p < 0.001$).

Conclusion: This study confirms the effectiveness of HBM-based nutritional education on healthy nutritional perception, belief and behavior among elderly women.

Copyright © 2017, Taiwan Society of Geriatric Emergency & Critical Care Medicine. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Population ageing has become a challenge for both developed and developing countries¹. The rapid changes in the number of aged people create important concern, particularly in relation to the health of seniors². Ageing is associated with reduction in both physiological and psychological function³ that may lead to a significant level of dependency⁴. The incidence of age-related diseases has largely increased over time⁵. So, most patients admitted into the hospitals are older adults and two-thirds of them have two or more chronic illnesses⁶.

The adoption of lifestyle behaviors such as smoking, poor nutrition, physical inactivity has resulted in higher levels of risk

factors for non-communicable diseases (NCDs).⁷ Elderly population are more prone to NCDs.⁸ Studies showed that these diseases highly overcome in Iranian elderly⁹ and its prevalence in elderly women are more than men¹⁰. Burden of NCDs and the role of socio-demographic and behavioral predictors among these group of population need to be investigated in detail⁸.

The hazardous effects of behavioral and nutritional risk factors on NCDs have been established in the cohort and randomized trials studies^{11,12}. Health educational program can help older people to adopt healthy lifestyle behaviors and prevent and control age-related diseases¹³. To change the behavior, however, it is essential to understand and apply the widely used models to predict healthy behavior. Older people have limited knowledge about health threatening behaviors¹⁴. In this line, various studies have used the educational models such as Health Belief Model (HBM), planned behavior and social cognition theory¹⁵. HBM model was applied much more than other models for behavioral changes and

* Corresponding author.

E-mail address: Jamileh.Amirzadeh@gmail.com (J. Amirzadeh Iranagh).<http://dx.doi.org/10.1016/j.ijge.2017.08.007>1873-9598/Copyright © 2017, Taiwan Society of Geriatric Emergency & Critical Care Medicine. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

prevention of diseases^{16,17}. Hence, the HBM was used as the framework of this study to explain effective factors to change nutritional behavior (perceived severity, perceived susceptibility, perceived benefit, perceived barrier, and self-efficacy) of the elderly women.

2. Material and methods

This study was approved by the Ethics Committee of Urmia medical Science University (P6/92/4/47647) as well as Universiti Putra Malaysia [UPM/TNCPI/RMC/1.4.18.1 (JKEUPM)/F1].

The sample of this randomized controlled trial study was 100 elderly women who attended primary health centers in the Urmia city from October 2013 to May 2014. This study included women aged 60 years and older who have been resident of Urmia city at least for 5 years and were independent in basic activities of daily living. The volunteers with NCDs diagnosis or <6 scores on the abbreviated mental test scale were excluded from this study.

The elderly women were recruited using multi-stage cluster random sampling technique. First, Urmia was divided into four districts. Then, one health center was selected randomly from each district. A total of 200 elderly women were randomly selected from the list of all households under the cover of these four health care centers. A number of 100 out of 200 elderly women were qualified and interested to participate in this randomized controlled trial study. They were allocated into intervention ($n = 50$) or control ($n = 50$) groups randomly. The written consent was signed by the participants after informing about the aim, benefits and drawbacks of the study.

The following equation was used for determining the sample size.

$$n = \frac{(\sigma_1^2 + \sigma_2^2/\kappa)(Z_{1-\alpha/2} + Z_{1-\beta})^2}{\Delta^2}$$

$n1$ = sample size of Group 1, $n2$ = sample size of Group 2, $\sigma1$ = standard deviation of Group 1. $\sigma2$ = standard deviation of Group 2, Δ = difference in group means, κ = ratio = $n2/n1$. $Z_{1-\alpha/2}$ = two-sided Z value, $Z_{1-\beta}$ = power.

The sample size was estimated 38 subjects in intervention and 38 for control group. However, considering a 30% possible attrition rate, 50 participants were finalized for each group.

2.1. Data collection

Data were collected using a self-administered and validated questionnaire which was prepared for evaluation the nutrition behavioral change based on Health Belief Model among Iranian elderly^{18,19}. The questionnaire consisted of four parts. The first part was demographic characteristics such as age, marital status, employment, educational level, housing status and living arrangement. The second part was knowledge of healthy nutrition that was measured by 10 items. The third part consisted of 21 items to evaluate the respondent's nutrition behavior for all of food groups. Each item measured the number of servings of the various food per day. Each correct response was scored one point and a wrong response was given zero point. Then, the total score was computed. Finally, beliefs of the participants were measured using (24) items. A 4-point Likert scale, ranging from 0 to 4 (0 = completely disagree, 1 = disagree, 2 = I don't know, 3 = agree, and 4 = completely agree) was used to measure the beliefs including perceived susceptibility (4), perceived severity (5), perceived benefits (5), perceived barriers (5), perceived self-efficacy (5). The reliability of this questionnaire was checked and

Table 1
Result of reliability test on HBM scale for the nutrition questionnaire.

Variable	Number of items	Spearman's correlation	Cronbach's alpha
Nutritional knowledge	10	0.980	
Nutritional behavior	21	0.868	
Perceived benefit	5		0.894
Perceived barriers	5		0.827
Perceived severity	5		0.895
Perceived susceptibility	4		0.798
Self-efficacy	5		0.920

reported in Table 1. The questionnaire was completed by both groups before, at 3 and 6 months after the intervention program.

2.2. Intervention

The intervention group were trained in 24 sessions of 120 min each. Training sessions were conducted in groups of individuals and consisted of lecture, focus group discussion, question and answer, and showing video. The intervention group was provided pamphlets and educational booklet which were designed base on HBM to review the provided topics and to ensure educational continuity.

The intervention program was designed based on the components of the HBM for nutrition behavior improvement purpose. The program provided a blueprint for CVD prevention among elderly women. Therefore, the consideration of the program was improving the nutritional status and creating positive perception's changes of the elderly women.

The topics of the program's contents consisted of the importance of consumption of dairy products, meat and meat substitutes, fruits and vegetables, water and drinks, and reduction of consumption of salt for the elderly health promotion. Furthermore, the healthy nutritional behavior including non-using solid oils and using the alternative methods of cooking (steaming, boiling, and grilling) were explained. The recommendation of the lifestyle guidelines for Iranian elderly was used to instruct the subjects about consequences of unhealthy nutrition.

During the study, no educational program was implemented for the elderly women in the control group. However, for ethical consideration, the same educational program was provided for the control group at the end of the intervention program.

2.3. Data analysis

Data were analyzed using SPSS version 20 (SPSS® IBM, New York, U.S. A). Descriptive statistics were used to report the means (m) \pm standard deviations (SD) of the variable scores at three times measurements. Two way repeated measure ANOVA was used to detect mean differences between and within groups. The independent t -test was used to compare mean scores for the benefit, severity, susceptibility, and self-efficacy between the intervention and control groups before the intervention. All data were normally distributed using the Shapiro-Wilk test, Skewness, kurtosis, and histogram. Significance level was set at $\alpha < 0.05$.

3. Results

A total of 86 out of 100 elderly women (41 in the nutrition and 45 in the control group) completed six months' study program. Table 2 indicates the socio-demographic characteristics of the study subjects. As depicted in this Table, there was no significant

Table 2
Socio demographic characteristics of participants.

	Nutrition group		Control group		Test value	p-Value
	No	(%)	No	(%)		
Age (Mean \pm SD)	(70.54 \pm 5.80)		(70.09 \pm 5.06)		T = 2.239	0.086
Marital status						
Married	22	53.7	21	46.7	$\chi^2 = 1.375$	0.711
Widowed	19	46.3	24	53.3		
Occupation status						
Housewife	39	95.1	44	97.8	$\chi^2 = 7.284$	0.058
Retired	2	4.9	1	2.2		
Educational level						
literate	3	7.3	3	6.7	$\chi^2 = 4.083$	0.256
Illiterate	38	92.7	42	93.3		
Housing status						
Private	38	92.7	40	88.9	$\chi^2 = 9.822$	0.119
Rented	3	7.3	5	11.1		
Living arrangement						
Alone	11	26.8	13	28.9	$\chi^2 = 14.241$	0.281
With spouse only	13	31.7	16	35.6		
With spouse & children	8	19.5	4	8.9		
With children only	9	22	12	26.7		

Note: SD = Standard deviation, No = Number.

Table 3
Baseline knowledge, belief, and nutritional behavior of intervention and controls groups.

	Nutrition (n = 41)	Control (n = 45)	t	p Value
Nutritional knowledge	4.00 \pm 1.76	4.11 \pm 1.18	0.738	p = 0.530
Nutritional behavior	6.34 \pm 1.98	6.57 \pm 1.22	0.414	p = 0.743
Perceived benefit	2.97 \pm 0.70	3.00 \pm 0.71	0.213	p = 0.887
Perceived barrier	2.52 \pm 0.71	2.53 \pm 0.79	1.316	p = 0.271
Perceived severity	3.33 \pm 0.42	3.51 \pm 0.62	0.814	p = 0.488
Perceived susceptibility	3.37 \pm 0.43	3.43 \pm 0.64	0.520	p = 0.669
Self-efficacy	3.32 \pm 0.62	3.52 \pm 0.74	1.331	p = 0.266

difference between two groups at the baseline. It should be noted that no one of the participants was smoking or drinking alcohol.

Table 3 shows the participants' knowledge, behavior, and belief at the baseline. As depicted in this Table, there were no significant differences in scores of the dependent variables between intervention and control groups.

Compliance or attendance rate for intervention group was 82%. This high response rate might be related to regular contact and reminding of the subjects to join the intervention sessions. Another reason may be associated with the social aspect of this program and the subjects' satisfaction with the nutritional education.

3.1. Outcomes of intervention

Before the intervention, the elderly women showed low level of the nutritional knowledge. After the intervention, nutrition group showed a significant improvement on nutritional knowledge compared to baseline ($F = 168.544$, $p < 0.001$, partial $\eta^2 = 0.501$) (see Fig. 1).

The statistical analysis for nutritional behavior showed a significant main effect for time ($F = 153.727$, $p < 0.001$, partial $\eta^2 = 0.478$) and group ($F = 69.036$, $p < 0.001$, $\eta^2 = 0.552$) for nutritional group, indicating a significant improvement in nutritional behavior scores across the three measurements and compared to the control group (Fig. 2).

After intervention, perceived benefit increased significantly in the intervention group vs. control group (mean differences in the intervention vs. control group: 1.576 vs. 0.200 respectively, $p < 0.001$). The results were the same for perceived barrier (mean

differences in the intervention vs. control group: 1.663 vs. 0.129 respectively, $p < 0.001$), severity scores (mean differences in the intervention vs. control group: 1.39 vs. 0.156 respectively, $p < 0.001$), susceptibility (mean differences in the intervention vs. control group: 1.189 vs. 0.122 respectively, $p < 0.001$), and self-efficacy (mean differences in the intervention vs. control group: 1.449 vs. 0.022 respectively, $p < 0.001$). However, perceived barrier scores decreased significantly in the intervention group compared to the control group (mean differences in the intervention vs. control group: 1.663 vs. 0.129 respectively, $p < 0.001$). Furthermore, all the mentioned variables changed significantly from pre-to post-test 1 and 2 and from post-test 1 to 2 only in the intervention group ($p < 0.001$, Table 4).

4. Discussion

The results of the present study showed the efficacy of a HBM-based nutritional education on the elderly women's knowledge, belief and behavior. Before the intervention, mean of nutritional knowledge for both groups was fairly poor. The nutritional knowledge grades in the previous studies were also in the low levels^{20,21}. After the intervention, the results of the present study are in accordance with findings of some studies^{22,23} that found a significant improvement in participants' knowledge after a nutritional education program. The method of training including face to face discussion, interactive lessons, showing video, group presentations, and question and answer sessions may be the key elements of effectiveness of the intervention used in this study²⁴.

After the intervention, in the present study, the mean values of the HBM components including perceived susceptibility, severity, benefits increased and perceived barriers decreased significantly in the intervention group compared to the controls. Similarly, researchers²⁰ applied a HBM based education intervention and observed a significant improvement in perceived susceptibility, severity, benefits and barriers for calcium intake. Likewise, another in Iran²⁵ stated that the educational intervention program had a significant effect on HBM constructs after 3 and 6 months intervention. One study²⁶ also showed that perceived susceptibility, benefit, and severity significantly increased and perceived barrier decreased in the nutritional education group. These results may be caused by the effectiveness of HBM-based educational program on improvement of participants' perception about healthy behavior.




Fig. 1. Nutritional knowledge of intervention and controls groups during intervention.




Fig. 2. Nutritional behavior of intervention and controls groups during intervention.

The results of this study are in agreement with the findings of the previous studies which indicated that the interventions positively influenced nutritional behavior among healthy^{18,27–31}. Although some others have reported in unhealthy elderly^{31–35}. An author²⁵ reported that the HBM-based nutritional education is more effective in positive behavioral changes compared to usual educational program (a lecture approach). This means that intervention group felt the hazards of poor adherence (the increase of perceived susceptibility) and the benefits of high adherence (the increase of perceived benefits) to the healthy diet that encouraged them to follow the healthy diet behaviors²⁹.

One study³⁶ although found a significant increase in the perceived susceptibility, benefits, and barriers of the healthy diet among the elderly residing in nursing home, but they could not observe a significant improvement in their nutritional behavior. Furthermore, another in Sweden³⁷ failed to detect nutritional behavior changes among 96 community-dwelling frail elderly

women after the 12-week nutritional intervention program and a six-month period follow up. This non-significant results have possibly been related to source of the sample and the lack of supervision through the intervention period.

The results of this study are in agreement with the findings of the previous studies which have supported the incorporation of self-efficacy into HBM based educational program. Self-efficacy is an effective factor for encouraging to follow the health behavior^{28,38,39}. For instance, a study⁴⁰ reported that weight loss intervention improved self-efficacy for the self-regulation of nutritional behavior among elderly people which influenced on their weight loss.

5. Conclusion

In conclusion, HBM-based education improved the knowledge, belief and behavior of the elderly. Therefore, the results of the

Table 4
Repeated measures ANOVA of belief of nutrition within groups.

	Time (I)	Time(J)	Nutrition group		Control group	
			Mean diff (J-I)	p value	Mean diff (J-I)	p value
Perceived Benefit	pre-test	postest1	1.307	0.001*	0.187	0.064
	pre-test	postest2	1.576	0.001*	0.200	0.061
	postest1	postest2	0.268	0.001*	0.013	0.824
Perceived Barrier	pre-test	postest1	0.956	0.001*	0.124	0.418
	pre-test	postest2	1.663	0.001*	0.129	0.380
	postest1	Postest2	0.707	0.001*	0.004	0.969
Perceived Severity	pre-test	postest1	0.976	0.001*	0.147	0.069
	pre-test	postest2	1.390	0.001*	0.156	0.060
	postest1	postest2	0.415	0.001*	0.009	0.877
Perceived Susceptibility	pre-test	postest1	0.860	0.001*	0.128	0.118
	pre-test	postest2	1.189	0.001*	0.122	0.13
	postes1	postest2	0.329	0.001*	0.006	0.937
Perceived Self-efficacy	pre-test	postest1	0.985	0.001*	0.004	0.965
	pre-test	postest2	1.449	0.001*	0.022	0.839
	postest1	postest2	0.463	0.001*	0.018	0.818

Note: Mean Diff = Mean Difference, * = $p < 0.001$.

present study support the efficacy of the HBM-based nutritional education among the community dwelling elderly people.

Conflicts of interest

None.

Acknowledgement

Thanks to the Universiti Putra Malaysia and its lecturers for careful and timely supervision throughout this PhD thesis. We thank all elderly woman for their participation in this study.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.ijge.2017.08.007>.

References

- Ella NARA, Shehab DI, Ismail MA. Prevalence of overweight and obesity, and status of chronic non-communicable diseases and some related risk factors among Egyptian adolescents. *J Diabetes Endocrinol.* 2011;2:41–52.
- Thanakwang K, Soonthornhadha K, Mongkolprasoe J. Perspectives on healthy aging among Thai elderly: a qualitative study. *Nurs Health Sci.* 2012;14:472–479.
- Mazzonna F, Peracchi F. Ageing, cognitive abilities and retirement. *Eur Econ Rev.* 2012;56:691–710.
- Singh GK, Hiatt RA. Trends and disparities in socioeconomic and behavioural characteristics, life expectancy, and cause-specific mortality of native-born and foreign-born populations in the United States, 1979–2003. *Int J Epidemiol.* 2006;35:903–919.
- Christensen K, Doblhammer G, Rau R, et al. Ageing populations: the challenges ahead. *Lancet.* 2009;374:1196–1208.
- Duque S, Freitas P, Silvestre J, et al. Prognostic factors of elderly patients admitted in a medical intermediate care unit. *Eur Geriatr Med.* 2011;2:327–331.
- Jemal A, Bray F, Center MM, et al. Global cancer statistics. *CA Cancer J Clin.* 2011;61:69–90.
- Feng L, Li P, Wang X, et al. Distribution and determinants of non communicable diseases among elderly Uyghur ethnic group in Xinjiang, China. *PLoS One.* 2014;9:e105536.
- Tanjani PT, Motlagh ME, Nazar MM, et al. The health status of the elderly population of Iran in 2012. *Arch Gerontol Geriatr.* 2015;60:281–287.
- Almasi Hashyani A, Salehi M, Torke-Jokar B, et al. Evaluation of nutritional behavior of elderly people about prevention of gastric ulcers in Shiraz. *Daneshvar.* 2013;20:43–50.
- Ezzati M, Riboli E. Behavioral and dietary risk factors for noncommunicable diseases. *N Engl J Med.* 2013;369:954–964.
- Lima-Costa MF, Matos DL, Ribeiro ALP. Chagas Disease predicts 10-year stroke mortality in community-dwelling elderly the Bambuí cohort study of aging. *Stroke.* 2010;41:2477–2482.
- Olsen JM, Nesbitt BJ. Health coaching to improve healthy lifestyle behaviors: an integrative review. *Am J Health Promot.* 2010;25:e1–e12.
- Dobe M. Health promotion for prevention and control of non-communicable diseases: unfinished agenda. *Indian J Public Health.* 2012;56:180.
- Prestwich A, Sniehotta FF, Whittington C, et al. Does theory influence the effectiveness of health behavior interventions? Meta-analysis. *Health Psychol.* 2014;33:465.
- Amirzadeh Iranagh J, Rahman HA, Motalebi SA. Health Belief Model-based intervention to improve nutritional behavior among elderly women. *Nutr Res Pract.* 2016;10(3):352–358.
- Corace KM, Srigley JA, Hargadon DP, et al. Using behavior change frameworks to improve healthcare worker influenza vaccination rates: a systematic review. *Vaccine.* 2016;34:3235–3242.
- Davari S, Dolatian M, Maracy MM, et al. The Effect of a Health Belief Model (HBM)- based educational program on the nutritional behavior of menopausal women in Isfahan. *IJME.* 2011;10:1263–1272.
- Noroozi E, Kasiri- Dolatabadi N, Eslami A, et al. Knowledge and attitude toward menopause phenomenon among women aged 40–45 years. *J Educ Health Promot.* 2013;2:1–5.
- Mohamed RA, Awad MM, Shalaby SI, et al. Effect of nutritional health education program on elderly nutritional knowledge, attitude and practice in Abu Khalifa primary health care center, Ismailia Governorate. *Med J Cairo Univ.* 2013;81:405–409.
- Shahar S, Adznam SN, Rahman SA, et al. Development and analysis of acceptance of a nutrition education package among a rural elderly population: an action research study. *BMC Geriatr.* 2012;12:1–9.
- Shojaeizadeh D, Sadeghi R, Tarrahi MJ, et al. The effect of educational intervention on prevention of osteoporosis through Health Belief Model (HBM) in volunteers of Khorramabad city's health centers in 2010–2011. *Ann Biol Res.* 2012;3:300–307.
- Sharifrad G, Tol A, Mohebi S, et al. The effectiveness of nutrition education program based on health belief model compared with traditional training. *J Educ Health Promot.* 2013;2:1–15.
- Kupka-Schutt L, Mitchell ME. Positive effect of a nutrition instruction model on the dietary behavior of a selected group of elderly. *J Nutr Elder.* 1992;12:29–53.
- Bayat F, Shojaezadeh D, Baikpour M, et al. The effects of education based on extended health belief model in type 2 diabetic patients: a randomized controlled trial. *J Diabetes Metab Disord.* 2013;12:45.
- Sharifrad G, Entezari M, Kamran A, et al. The effectiveness of nutritional education on the knowledge of diabetic patients using the health belief model. *JRMS.* 2009;14:1–6.
- Gutiérrez-Doña B, Lippke S, Renner B, et al. Self-efficacy and planning predict dietary behaviors in Costa Rican and South Korean women: two moderated mediation analyses. *Appl Psychol Health Well-Being.* 2009;1:91–104.
- Hamuleh M, Vahed S, Piri AR. Effects of education based on Health Belief Model on dietary adherence in diabetic patients. *J Diabetes Metab Disord.* 2010;9:1–6.
- Bhuroo T, Jeewon R. Effectiveness of a theory-driven nutritional education program in improving calcium intake among older mauritian adults. *Sci World J.* 2013;2013:1–17.
- Nadine RS. Nutrition education for the healthy elderly population: isn't it time? *J Nutr Educ Behav.* 2002;34(Suppl 1):S42–S47.
- Wells JL, Dumbrell AC. Nutrition and aging: assessment and treatment of compromised nutritional status in frail elderly patients. *Clin Interv Aging.* 2006;1:67–79.

32. Payette H, Boutier V, Coulombe C, et al. Benefits of nutritional Supplementation in free-living, frail, undernourished elderly people: a prospective randomized community trial. *J Am Diet Assoc.* 2002;102:1088–1095.
33. Gaines JM, Narrett M, Parrish JM. The effect of the addition of osteoporosis education to a bone health screening program for older adults. *Geriatr Nurs.* 2010;31:348–360.
34. DeNysschen CA, Brown JK, Cho MH, et al. Nutritional symptom and body composition outcomes of aerobic exercise in women with breast cancer. *Clin Nurs Res.* 2011;20:29–46.
35. Pinto SL, Lively BT, Siganga W, et al. Using the Health Belief Model to test factors affecting patient retention in diabetes-related pharmaceutical care services. *Res Soc Adm Pharm.* 2006;2:38–58.
36. Mamaghani ME, Mahdavi R, Vaziri Y. Dental health and its relation to macro-nutrient intake and anthropometric indices "khuban" private nursing home residents in Tabriz. *J Tabriz Univ Med Sci.* 2007;29:21–27.
37. Rydwick E, Frandin K, Akner G. "Effects of a physical training and nutritional intervention program in frail elderly people regarding habitual physical activity level and activities of daily living" A randomized controlled pilot study. *Arch Gerontol Geriatr.* 2010;51:283–289.
38. Jeffery RW, Bjornson-Benson WM, Rosenthal BS, et al. Correlates of weight loss and its maintenance over two years of follow-up among middle-aged men. *Prev Med.* 1984;13:155–168.
39. Swaim RA, Barner JC, Brown CM. The relationship of calcium intake and exercise to osteoporosis health beliefs in postmenopausal women. *Res Soc Adm Pharm.* 2008;4:153–163.
40. Rejeski WJ, Mihalko SL, Ambrosius WT, et al. Weight loss and self-regulatory eating efficacy in older adults: the cooperative lifestyle intervention program. *J Gerontol B Psychol Sci Soc Sci.* 2011;66:279–286.