

Towards Realizing Benefits of Information Technology in Organ Transplant: A Review

Parviz RASHIDI KHAZAE^a, Habibollah PIRNEJAD^{b,c}, Jamshid BAGHERZADEH^a,
and Zahra NIAZKHANI^{b,d,1}

^a*Electrical and Computer Engineering Department, Urmia University*

^b*Department of HIT, Urmia University of Medical Sciences, Urmia, Iran*

^c*Institute of Health Policy and Management (iBMG), Erasmus University Rotterdam,
Rotterdam, The Netherlands*

^d*Nephrology and Kidney Transplant Research Center, Urmia University of Medical
Sciences, Urmia, Iran*

Abstract. Organ transplantation comprises of many phases, processes, and activities and involves multiple stakeholders. Effective management of such a complex and costly medical domain requires an efficient, multifaceted solution. Although, Information Technology (IT) can basically play an important role here, it is not clear how IT potentials have been deployed so far. We systematically reviewed MEDLINE, EMBASE, CINAHL, The Cochrane and IEEE databases and identified 27 publications describing IT application in organ transplantation. Although the IT coverage spans over waiting list management, donor-recipient matching, and inpatient and outpatient medication and lab monitoring practices, the coverage is still patchy and whole process IT support is missing in practice.

Keywords. Organ Transplantation, Information Technology, systematic review

Introduction

Replacing a defective organ with a healthy one has opened a new horizon in medical practice. With advances in transplantation filed over the last fifty years, many technical and medical hurdles have been overcome; and as a result of this, patients' survivals and their quality of life have greatly been improved [1]. Yet, there are many challenges to be met. Transplantation process, from organ procurement to recipients' follow ups, can practically be divided into 3 main interdependent phases of pre-transplant, inpatient, and follow-up care involving different stakeholders such as clinicians, provider organizations, donors and recipients. Information sharing and coordination among these stakeholders are at the center stage of the whole process; and if not managed properly, the quality and efficiency of the process can be hampered. In this regard, Information Technology (IT) has great potential to offer by removing many hurdles in the processes of this very specialized field. However, it is not clear how far IT potentials have been deployed and which areas still need more consideration.

¹ Corresponding Author. Dr. Zahra Niazkhani, Urmia University of Medical Sciences, Postal Code: 5714783734, Urmia, Iran. E-mail: niazkhani.z@umsu.ac.ir.

1. Methods

A systematic search was conducted in main databases (details in Figure 1, last search on Dec. 4th, 2015). Through a Boolean search strategy using key and MeSH terms related to *transplantation AND IT intervention*, we identified English qualitative or quantitative studies that described IT use by *care providers* to support organ transplantation’s daily inpatient and/or outpatient clinical workflow. Editorials, opinion papers, reviews, stem cell transplant literature, simulation studies, and literature lacking a routinely used IT system in clinical workflow e.g., registries or systems used merely for collecting data for research purposes were excluded. We analyzed and categorized the final set of literature on the bases of the processes supported by IT systems.




Figure 1. Flow diagram of study selection (search from Jan. 1990 to Dec. 2015)

2. Results

Twenty-seven publications describing IT application in each phase of organ transplant care were identified. These systems’ main application area are described below and summarized in Table 1. The Pre-Transplant Phase: Managing rapidly increasing number of patients waiting for transplantation through IT systems brought about close cooperation among transplant programs and allowed retrieving, integrating, analyzing, and interpreting numerous pieces of patient clinical information quickly [2-5]. IT systems have enabled local or national organ procurement organizations to manage shortages of organs more efficiently [6]. When an organ is available, matching and selection of the best recipient becomes an essential task. In this regard, IT systems such as telemedicine applications, PDAs, and online sharing of information have successfully been used to facilitate the process with the aim of improving outcome, quality of life and longevity

Table 1. Summary of the literature describing the coverage of different processes in transplantation care by IT systems.

Phases	Application areas	References
Pre-Transplant	Waiting List	[2, 5, 7, 8, 10]
	Organ Procurement	[6]
	Matching best candidate	[3, 7, 8, 9]
	Decision Making	[3, 4]
	Survival Analysis	[7]
Inpatient	Med. CPOE/CDSS	[13-16]
	Lab CDSS	[17]
	Barcode-assisted medication administration	[18]
	Patient Education	[19]
Post-transplant Follow up	Outpatient Lab values Monitoring	[20-23]
	Immunosuppressive & other Med Monitoring	[22, 24-27]
General*	Managing clinical information in diff. phases	[5, 9, 11, 12, 28]

* These IT applications supported activities of more than one phase of organ transplant.

of patients [3, 4, 7-10]. Before the hospitalization of donors and recipients, IT has also been used in supporting clinicians to manage and to evaluate information coming from different consultant physicians, and the results of para-clinical and laboratory investigations [5, 9, 11, 12].

The Inpatient Transplant Phase: Literature also described IT systems used in inpatient settings for better caring of grafted organs and for preventing its rejection or failure. These systems mainly focused on managing dosages of immunosuppressive and other medications, for example through applying CPOE and reducing incidence or severity of potential drug-drug interactions [13-16]. One study also explored the benefits of IT in managing lab resource utilization [17] and another on safety of inpatient medication process via application of Barcode-assisted Medication Administration System [18]. Another study reported on the application of an IT system to promote evidence-based inpatient education for improving the result of transplantation [19].

The Post Transplant Follow-up Phase: Post-transplantation care and lifelong follow-ups are required to protect patients against transplant rejection and also to detect and manage complications of long-term immunosuppressive drugs such as cancer and diabetes. Close monitoring of Lab values is a key to detect the early signs of graft rejection. IT systems have successfully been used to help clinicians to monitor relevant Lab values [20-23]. These systems can provide an integrated view of patient data over time and notify clinicians whenever changes in Lab values require special attention [23]. Literature reported application of IT systems and Informatics methods in order to reduce toxicity of immunosuppressive drugs (e.g., Tacrolimus) while making sure that patients receive necessary dosage [24-26]. IT systems have also supported clinicians in monitoring other medications such as antiviral drugs or anti-lipid therapy whenever needed [20, 27].

3. Discussion and Conclusion

Because of the existing need, many endeavors have been taken place towards the application of IT systems in organ transplantation. However, our review shows that the coverage is still patchy with a greater emphasis on pre-transplant process. This is most probably because managing of scarce resources of donated organs efficiently is of great importance. In fact, despite a comprehensive search, we could not find any setting that has had utilized a comprehensive IT solution that provides support for the all phases of organ transplantation. Given the fact that preserving the transplanted organ is also of great importance, IT solutions are also required to support lifelong post-transplant care. Supporting the implementation of well-established but complex, standard care protocols pertaining to transplantation would be a considerable value added by IT systems.

References

- [1] C.J. Watson, J.H. Dark, Organ transplantation: historical perspective and current practice, *Br J Anaesth* **108** (Suppl 1) (2012), i29-42.
- [2] H.Y. Bircan, Ü. Özcelik, B. Koc, M.E. Aras, D. Okutur, A. Demiral, Development of an information model for kidney transplant waiting list, *Exp Clin Transplant* **13** (Suppl 3) (2015), 58-61.
- [3] B. Barshick, How PDAs can speed up the organ transplant process, *Nursing* **34** (4) 2004, 68.

- [4] A. Krause, D. Hartl, F. Theis, M. Stangl, K.E. Gerauer, A.T. Mehlhorn, Mobile decision support for transplantation patient data, *Int J Med Inform* **73** (5) (2004), 461-464.
- [5] M. Kurtz, T. Bennet, et al., Demonstration of SLUMIS: a clinical database and management information system for a multi organ transplant program, *Proc Annu Symp Comput Appl Med Care* (1992), 889-890.
- [6] P. López-Álvarez, F. Caballero, S. Willmott, U. Cortés, A. López-Navidad, CARREL: An internet platform for the distribution of human organs for transplantation, *Transpl Proc* **37** (9) 2005, 3667-3668.
- [7] S.G. Vari, G. Brugal, F. Godo, B. Bercic, G. Nagy, G. Avar, et al., Regional and international integrated telemedicine network for organ transplant, *Proceedings / AMIA 2000; Annual Symposium*, 873-877.
- [8] J.C. Ruiz, R. Escallada, J.G. Cotruello, J.A. Zubimendi, et al., Model of a telematic network for communication between centers in a kidney transplant area, *Transplant Proc* **31** (6) 1999, 2358.
- [9] S.H. Mitchell, P. Zaltash, A.L. Miller, R.D. Gordon, Transplant information management system: a center-oriented approach to transplant data management, *Top Health Rec Manage* **11** (2) 1990, 20-24.
- [10] C. Wu, N. Shah, P. Sood, C. Puttarajappa, J. Bernardo, R. Mehta, A. Tevar, R. Shapiro, H. Tan, M. Wijkstrom, M. Sturdevant, S. Hariharan, Use of the electronic health record (EHR) to improve the pre-transplant process for kidney and pancreas transplantation, *Transplantation* **98** 2014, 833-834.
- [11] K. Kirchner, C. Malessa, N. Herzberg, S. Krumnow, O. Habrecht, H. Scheuerlein, et al., Supporting liver transplantation by clinical pathway intelligence, *Transplant Proc* **45** (5) 2013, 1981-1982.
- [12] L.P. Dang, S. Ireland, N. Zuhdi, Modular Integrated Transplant Information System (MITIS), *Proc Annu Symp Comput Appl Med Care* (1994), 995.
- [13] K. Marfo, D. Garcia, S. Khalique, K. Berger, A. Lu, Evaluation of medication errors via a CPOE system in an inpatient renal transplant unit, *Transplant Research and Risk Management* **3** (2011), 91-96.
- [14] H.S. Mekhjian, R.R. Kumar, L. Kuehn, T.D. Bentley, P. Teater, A. Thomas, B. Payne, A. Ahmad, Immediate benefits realized following implementation of physician order entry at an academic medical center, *J Am Med Inform Assoc* **9** (5) 2002, 529-539.
- [15] P. Polidori, C. Di Giorgio, A. Provenzani, Incidence of potential drug interactions in a transplant centre setting and relevance of electronic alerts for clinical practice support, *Inf Prim Care* **20** (4) 2012, 257-262.
- [16] R. Tiwari, D.S. Tsapepas, J.T. Powell, S.T. Martin, Enhancements in healthcare information technology systems: customizing vendor-supplied clinical decision support for a high-risk patient population, *JAMIA* **20** (2) (2013), 377-380.
- [17] L. Boon-Falleur, E. Sokal, M. Peters, J.M. Ketelslegers, A rule-based decision support application for laboratory investigations management, *Proc Annu Symp Comput Appl Med Care* (1995), 314-318.
- [18] J. Bonkowski, R.J. Weber, J. Melucci, T. Pesavento, M. Henry, S. Moffatt-Bruce, Improving medication administration safety in solid organ transplant patients through barcode-assisted medication administration, *Am J Med Qual* **29** (3) (2014), 236-241.
- [19] V. Schaevers, A. Schoonis, et al., Implementing a standardized, evidence-based education program using the patient's electronic file for lung transplant recipients, *Prog Transplant* **22** (3) (2012), 264-270.
- [20] D.K. Hooper, C.L. Kirby, P.A. Margolis, J. Goebel, Reliable individualized monitoring improves cholesterol control in kidney transplant recipients, *Pediatrics* **131** (4) (2013), e1271-e9.
- [21] A. Sarkar, Supporting Longitudinal Care for Transplant Patients with an External Laboratory Data Entry Application, *AMIA Annual Proceedings* (2001), 1021.
- [22] N. Shah, C. Wu, et al., Improvement in quality of care metrics through the implementation of electronic health records (EHR) in renal transplant patient management, *Transplantation* **98** (2014), 814-815.
- [23] C.J. Staes, R.S. Evans, B.H.S.C. Rocha, et al., Computerized Alerts Improve Outpatient Laboratory Monitoring of Transplant Patients, *J Am Med Inform Assoc* **15** (3) 2008, 324-332.
- [24] A. Asberg, P. Falck, L.H. Undset, C. Dorje, H. Holdaas, A. Hartmann, K. Midtvedt, Computer-assisted cyclosporine dosing performs better than traditional dosing in renal transplant recipients: results of a pilot study, *Ther Drug Monit* **32** (2) 2010, 152-158.
- [25] E.S. Park, M.R. Peccoud, et al., Use of an automated clinical management system improves outpatient immunosuppressive care following liver transplantation, *J Am Med Inform Assoc* **17** (4) (2010), 396-402.
- [26] J.E. Ray, A.M. Keogh, A.J. McLachlan, Decision support tool to individualize cyclosporine dose in stable, long-term heart transplant recipients receiving metabolic inhibitors: overcoming limitations of cyclosporine C2 monitoring, *J Heart Lung Transplant* **25** (10) 2006, 1223-1239.
- [27] D. Hensler, C. Richardson, J. Brown, C. Tseng, P. DeCamp, M. Ison, Effect of an automated electronic medical record-based, pharmacist-driven valganciclovir dose optimization program in solid organ transplant recipients, *Transplantation* **98** (2014), 763.
- [28] R.M. Merion, A.B. Leichtman, N. Tsui, J. Dean, Iterative cooperative prototyping in the design of web-based transplant information systems, *Transplant Proc* **30** (5) (1998), 1634-1636.