

# Pharmaceutical, phytochemical, and economical potentials of *Glycyrrhiza glabra* L: a review

Mahmoud Bahmani<sup>1</sup>, Amir Sarrafchi<sup>2</sup>, Hedayatollah Shirzad<sup>2</sup>, Najmeh Shahinfard<sup>2</sup>, Mahmoud Rafieian-Kopaei<sup>2\*</sup>, Somayeh Shahsavari<sup>3</sup>, Babak Baharvand-Ahmadi<sup>4</sup>, Morovat Taherikalani<sup>4</sup>, Sobhan Ghafourian<sup>3</sup>

<sup>1</sup>Food and Beverages Safety Research Center, Urmia University of Medical Sciences, Urmia, Iran

<sup>2</sup>Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

<sup>3</sup>Clinical Microbiology Research Center, Ilam University of Medical Sciences, Ilam, Iran

<sup>4</sup>Razi Herbal Medicines Research Center, Lorestan University of Medical Sciences, Khorramabad, Iran

\*Corresponding author: E-Mail: rafieian@yahoo.com

## ABSTRACT

*Glycyrrhiza glabra* L. (Licorice) is one of the most important medicinal plants indigenous to Iran. The main component of this species is a triterpenoid saponin compound called glycyrrhizic acid or glycyrrhizin that is 30-40 times sweeter than sucrose and is used in medicine, food and tobacco industries. Licorice has different therapeutic implications as it is used against Hepatitis c, skin and pulmonary diseases, and heart failures. It also improves the immune system and has anti-inflammatory, antiviral, antimicrobial, antioxidant and anticancer properties. The aim of the present study is to demonstrate the pharmaceutical, phytochemical, and economic potentials of this plant.

**KEY WORDS:** Therapeutic effects, Licorice, *Glycyrrhiza glabra*, Glycyrrhizic acid, Glycyrrhizin.

## 1. INTRODUCTION

It has been estimated that the burden of hard curable diseases including diabetes mellitus, atherosclerosis and other cardiovascular diseases, related to oxidative stress, contribute for about 60% of the 56.5 million reported deaths in 2001 (Farnsworth, 2004). Nowadays, a lot of researches focus on identification and preparation of natural compounds for prevention (Bahmani, 2014; Rahnama, 2014; Rafieian-Kopaei, 2014; Delfan, 2014; Asgary, 2013) and treatment (Rafieian-Kopaei, 2014; Gharipour, 2013; Asadi-Samani, 2014; Mirhosseini, 2014) of these diseases. Medicinal plants are a reliable source for preparation of effective drugs (Bahmani, 2014; Khosravi-Boroujeni, 2012; Saki, 2014; Khosravi-Boroujeni, 2013; Karamati, 2014).

Licorice with the scientific name of *Glycyrrhiza glabra* L. (Figure 1) is one of these plants. It is a perennial herb of fabaceae family which possesses a wide range of medicinal and food compounds in its root and rhizome. Licorice is widely used in medicinal, food and also tobacco industries. However, it grows easily so that it decreases products in farms and gardens due to high development of root and rhizome (Chandler, 2000). With regard to increasing value and special position of this plant in new medicinal industries, it is necessary to recognize its various potentials in different areas that is the aim of this review paper.



**Figure 1:** *Glycyrrhiza glabra*

**1.1. Plant name and botanical description:** The scientific name of licorice is *Glycyrrhiza glabra* L. and is known as Licorice or Liquorice. *Glycyrrhiza glabra* is a Greek name and has been derived from two words of glykys meaning sweet and rrhiza means root. Glabra means smooth refers to the fruit of this species which is smooth (Akhundzadeh, 2000).

Licorice is a perennial plant of the Astragaleae family and subfamily of Fabaceae and Faboideae. Height of this plant is different and reaches to 100-200 cm. The plant has mass and excessive leaves. The leaves are alternatively attached to the stem and have 4-7 leaflets and a terminal leaflet. Leaflets are dark green. Flowers are seen irregular and crowded on terminal spikes in yellow, violet or purple colors. Length of flowers reaches to more than 1 cm. Plant flowers late spring and early summer (June-July). Fruit is an oblong pod, 2-3cm long and brown in color. Sides of fruit are thin and become more or less peaked. There are 3-5 brown seeds inside the fruit

that are like bean. Seed layer is thick and steady and weight of one thousand seeds is about 10 gram. Length of root is different between 30-60 cm depending on species of plant and climatic condition of growing place. Length of root in dry regions and light soils reaches to 200 cm. Root and rhizome have medicinal uses. Licorice has different varieties including *G. glabra* var. *glandulifera*, *G. glabra* var., *G. glabra* var. *violacca* and *G. glabra* var. *pallid* (Ghahraman, 1999). *Glycyrrhiza glabra* grows in deep valleys with full sun mostly in well-drained soils (Merriam-Webster's, 2007).

**1.2. Distribution:** Licorice is a Mediterranean plant and is cultivated extremely in south-east of Asia. This plant grows from south of Europe to central Asia between two 30-45°N latitudes. The countries producing liquorice include Iran, Pakistan, Afghanistan, Iraq, China, Turkmenistan, Azerbaijan, Uzbekistan, and Turkey. Licorice is also cultivated in extensive areas in England, Belgium, French, Greece, Italy, and Germany. In Iran, also this plant is frequently found in whole east, west and center of country. Although in many countries this plant is cultivated as a medicinal plant but is considered as a weed in wheat and pea farms of central regions of Iran like Isfahan and Arak and dry farms of Kermanshah, Ilam and Fars provinces (Ghahraman, 1999).

**1.3. Commercial applications and preparation:** Licorice is the dried root of *Glycyrrhiza glabra* that has valuable chemicals such as glucose, sucrose, asparagine, albomidic substances, resin and essence. Effective substances of this plant have various consumptions in pharmacy, tobacco and confectionary industries (Mirhaidar, 1994; Fenwick, 1990). Licorice has a favorable flavor and taste. Its flavor mainly comes from *anethole*. This compound is also available in anise, fennel, and some other herbs. As a flavoring agent, licorice is used for chewing tobacco, liqueurs and soft drinks. The sweetness of licorice comes mainly from *glycyrrhizic acid* (Fenwick, 1990).

Licorice extract contains approximately 100 calories per ounce. By boiling root of plant and volatilizing its main part of water, a black substance is obtained. The most important indexes to determine quality of licorice root are the amount of aqueous extract and the amount of glycyrrhizic acid percentage. Amount of extract should be at least 20% in dried roots of plant 17 and the percentage of glycyrrhizic acid in plant root should be at least 2/5% according to USP pharmacopeia, 22 and 4 % (British Pharmacopeia, 2004) according to BP pharmacopeia. Free glycyrrhizin has been reported 58-60% and up to 98% as salt, in extract. In western countries, this substance is mainly used as sweetener since it is 50 times sweeter than sucrose and also has medicinal properties. Dried root of licorice can be chewed as sweet. Licorice extract is used in various types of sweet condiments (like candy). There are a lot of pharmaceutical products prepared from licorice, worldwide. Some of these products are used in Iran as medicine which are relatively popular and are sold as OTC (over the counter) are listed in table 1.

**Table 1: Medicinal products of licorice available in Iran**

Row	Drug Name	Medicinal form	Medicinal properties (manufacturer suggestion)	Name of manufacturer
1	Reglis	Pill	Treatment of duodenal and abdominal distension, excess acid secretion	Iran Darouk
2	d-reglis	pill	Inhibiting peptic ulcer in concurrent use with <i>Non-steroidal anti-inflammatory drugs</i>	Iran Darouk
3	Licophar	Pill	Anti-inflammatory, anti-cough and mucoactive	Goldaru
4	Gastrine	Pill	Anti-inflammatory, analgesic, accelerating duodenum and stomach ulcers healing	Goldaru
5	Mentazin	Pill	Relief of gastrointestinal pain, treat gastric ulcer and boating, laxative	Ebne-Masuyeh
6	Altadin	Pill	Treatment of inflammation and irritate mucous membranes, mucoactive in stimulated coughs	Dineh
7	Shirinoush	Syrup	Treatment of stomach and duodenum ulcer, gastritis and gastralogy	Goldaru
8	Reglicidin	Pill	Treatment of stomach and duodenum bloating, gastritis and gastralogy	Dineh

Some extracts of licorice are without glycyrrhizin and are called de-glycyrrhizinated licorice. This type of extract does not affect adrenal gland and is useful for people who suffer stomach and duodenum ulcers. This type of extract is used as basic substance in producing drinks.

#### 1.4. Plant cultivation:

**1.4.1. Ecological needs:** Origin of licorice is Mediterranean areas and needs hot weather and enough sunlight during growth. Licorice species grow in sandy-loam soil and bearing lime compounds. Root of this plant is extensively developed and its performance increase in sandy soils, rich of calcium and pH of 5/5-8/1. Enough

nutrients have main role in increasing root performance and its active ingredients. Licorice favors light and is well grown in 7-25 °C. Enough light, water and nutrients increase active ingredients of root. This plant usually grows in regions where annual rainfall varies from 400 to 1160 mm (British Pharmacopeia, 2004).


Early spring is suitable time to direct planting of grain in the main land 24. Early spring also is suitable time for planting seed in indirect method Suitable rows distance is 30-40cm. licorice seed should be planted to a depth of 2cm. The best time for vegetative reproduction of licorice is fall. Inter-row *spacing* is 60-80 cm and two shrubs are spaced 30-40cm apart in two rows. The suitable depth of planting root pieces is different and depends on soil texture and environment moisture so that the recommended depth in light soils is 15-20cm and 10-12 cm in heavy soils. Vegetative reproduction of licorice is more economical and often this method is used to reproduce this plant. Increasing planting depth from 20-52 cm increases leaf surface and then decrease it. Increasing planting depth and rhizome size increased dry weight. Dry weight of stem in low depths decreased but it increased when rhizome size increased. The highest total weight in planting depth obtained 20cm that was very close to dry weight of 40cm depth. Three-node Rhizome produced the highest dry weight than one-node and two-node rhizome (Waliullahpoor, 2005).

Licorice can be reproduced by seed or vegetative method. Since licorice seed has thick layer and this layer decreases its vegetative power, some scratch should be created in the surface of layer before planting it. Reproduction by seed is performed by two direct and indirect methods. Rhizomes do not germinate in temperature below 6 °C and need at least 15 °C for germination. This temperature has been considered as base temperature for germination of licorice (Nezamabadi, 2006).

**1.4.2. Maintenance and care:** Licorice growth in the first year is slow so that weeds can develop rapidly. It is necessary to keep weeds under control before planting and also preparing bed without weeds. In the first growth year, weeds should be controlled by hand or mechanically by cultivator. In the spring season in order to develop root, 10-cm long basal stems of licorice should be cut and bring out of land before leaves are fallen (Omidbeigi, 2004).

**1.4.3. Product harvest:** If licorice is reproduced by vegetative method, 3-4th year after planting its root can be harvested but if reproduction is done by seed its root can be harvested 5-6th year after planting (Ghanbari, 2005). The suitable time to use root is early fall to late winter with regard to specification of growth region that has the highest glycyrrhizin level in this time. To recovery root, furrows are made in land by tractor, axe or shovel at a depth of 50cm and roots are collected by worker. Since effective substance in licorice root are solved in root so roots should be washed rapidly and by flowing water. After cleaning, roots skin should be separated then divided into 10-15 cm pieces and finally dry them. To dry roots, direct light or high temperature should not be used. If electrical dryers are used, the suitable temperature is 40 °C. Dried roots are then cracked. Dried root performance in 3 year old plants reaches to 1.5-2 ton in hectare and ratio of fresh roots to dry roots is 3.5-1 24. It seems that roots with fewer diameters have more side surface for dry weight and can search soil and attract food elements better than thicker roots (Rengel and Graham, 1996). In roots with more diameters, skin and epidermis is destroyed in result of generative layer and one cork tissue is created (Mojtahedi and Lesani, 2015).


**1.5. Chemical compounds:** Licorice root contains a wide variety of compounds like sucrose (up to 18%), flavonoids, sterols, amino acids, starch, oil essences and saponins. The main triterpene is glycyrrhizic acid or glycyrrhizin (Figure 2) (C<sub>42</sub> H<sub>62</sub> O<sub>16</sub>) that is composed of two molecules of glucuronic acid and one molecule of glycyrrhetic acid (Jiang, 2004; Montoro, 2011; Blumenthal, 2000; Alan Teck, 2007). Glycyrrhizin salt can be as potassium or calcium (Sabbioni, 2005). This substance as the most important substance in licorice root is about 50 times sweeter than sugar (Hayashi, 1998). The amount of this substance in the root varies with plant variety and climatic conditions of growing place and is between 5-20%. For example, the amount of glycyrrhizic acid in Spanish licorice root is 6-8% and in Russian licorice is about 10-14%. Some researchers believe that Iranian licorice contains the highest level of glycyrrhizic acid (Douglas, 2004). Level of this substance increases following age increase so that root has the highest glycyrrhizic acid in final years. The sweetness of licorice is very different from sugar, being less instant, tart, and in comparison to sugar lasting longer (Somjen, 2000). Study of scientific resources on chemical and physical properties of this compound shows that glycyrrhizic acid decomposes in 120°C (Ong and Len, 2003). In some resources also softening point has been mentioned 170°C and melting point 220°C (Acharya, 1993). Different methods of extracting glycyrrhizic acid from root powder have been studied. Base of these methods often has been Pressurized hot water extraction (PHWE) and using alkalies and acids or both and other chemicals and adding them to hot water or steam to increase extraction use (Acharya, 1993). In Iranian National Standard and Pharmacopeia, the high-performance liquid chromatography has been used to determine glycyrrhizic acid level (Shi and Mei, 2003). Glycyrrhizic acid concentration in licorice root depends on variety, place and time of harvest (Hayashi, 1998).



**Figure 2. Structure of Glycyrrhizic acid**

Results of determining Glycyrrhizic acid and aqueous extract of samples collected from different regions of Iran have shown that plant grown in Kermanshah, Fars and Kerman border, has been the best product and has high export value. In addition to mentioned compounds, licorice root contains acid-2-beta-glycyrrhizic, glucuronic acid, glycyrrhetic acid (enoxolone), tannic acid, asparagine, resins, volatile oils, flavonoids such as liquiritigenin, liquiritin, isoliquiritigenin, and coumarin compounds, such as herniarin and umbelliferone with medicinal values (Li, 2000). Licorice also has phytoestrogens such as glabren and glabridin (Alan Teck, 2007).

Licorice root contains triterpene saponins, flavonoids, isoflavonoids and chalcones, coumarins, stilbenoids, as well as miscellaneous compounds as biologically active components. Most of them have antioxidant activities. The dehydroglyasperin D (DGD), isoangustone A (IsoA) and phenylflavonoids dehydroglyasperin C (DGC) have been derived from licorice and identified as major antioxidants of this plant (Figure 3) (Alan Teck, 2007). Among them DGC possesses the highest free radical scavenging capacity. All of these three compounds and the other components with antioxidant activity suppress lipid peroxidation in rat tissues as well as  $H_2O_2$ -induced free radical production in hepatoma cells (Hyo- Jung, 2012).



**Figure.3: Molecular structures of dehydroglyasperin C(A), dehydroglyasperin D(B), Isoangustone A(C).**

**1.6. Pharmacological and medicinal properties:** It is more than 4000 years that the root and rhizome of licorice has been used as medicine. It is recorded in pharmacopeia of different countries like America, China and other countries. In traditional medicine of Asia and Europe, Licorice is used to treat gastritis, respiratory infections and peptic complications (Bahmani, 2014). It is also used to treat hepatitis, tuberculosis, and other viral or microbial infections, as well as cancer, cough and heart diseases (Li, 2011). Glycyrrhizic acid has ability to inhibit *Helicobacter pylori* bacterium (Li, 2011) and so it is effective in the treatment of gastric ulcer and gastric mucosal problems (Csuk, 2010; Fukai, 2002). Licorice influences body endocrine gland system and its consumption can decrease testosterone level in blood. Also it has been demonstrated that Licorice is able to increase serotonin and prostaglandin secretion in stomach (Colalto, 2010). Commission E (The Complete German Commission E) has approved the use of licorice for gastric or duodenal ulcers as well as for catarrhs of the upper respiratory tract (Hayashi, 1998). Today also licorice extract is considered as one of ingredients of cough syrup (Hayashi, 1998). Extract of this plant also inhibits simulation of HIV virus in patients with AIDS (Alan Teck, 2007). Presence of lipid-lowering compounds and flavonoids with strong antioxidant activity and the positive effects of the extract on prevention of atherosclerosis have been approved in rabbit (Bahmani, 2014). However, studies have not yet reported that extract of this plant prevents and treats this disease in human (Fiore, 2008). Licorice has been shown promising effects on skin disease, especially against atopic dermatitis and hyperpigmentation as well as dental caries and neurodegenerative disorders (Gazzani, 2012; Messier, 2012).

**1.7. Side effects:** The United States FDA has announced that foods containing licorice or glycyrrhizin is safe if not consumed excessively. However, glycyrrhizin and enoxolone, the major components licorice inhibiting effects on cortisol degradation, Licorice may cause hypokalemia, hypertension, edema and weight gain or loss (Douglas,

2004; Somjen, 2000). No more than 100mg to 200mg of glycyrrhizin per day has been suggested, which is equivalent to about 70 to 150g of liquorice (Omar-Hesham, 2015).

Irregular and excessive consumption of licorice or other its products has been inhibited due to irritation of adrenal gland and excess secretion of aldosterone hormone. This state causes some effects like disorder in metabolic activities and high blood pressure. It has been reported that eating more than 20g per day or more probably causes undesirable effects. Excessive use of licorice is also harmful for spleen. Excessive use of licorice may lead to presence of hypertension and also heart stroke. Some people who use high doses of licorice feel myalgia and some other leg and hand swelling. High use of it increases weight (Maha, 2012).

Licorice use must be avoided in cases of hypertension or renal, heart and liver problems. Using this plant and its products has been inhibited for women in pregnancy or lactation periods. Of course these problems can be prevented if recommended doses of it are used. If long-term use of licorice is necessary, it is better to combine it with tragacanth (Maha, 2012).

## 2. DISCUSSION AND CONCLUSION

Licorice has been recognized as important medicinal plant through world. Different uses of this plant in pharmaceutical and food industries are reason of its significant commercial value in the world. Licorice possesses a wide variety of phytochemicals, having pharmacological and therapeutic properties. Licorice root contains triterpene saponins, flavonoids, isoflavonoids and chalcones, coumarins, stilbenoids, as well as miscellaneous compounds as biologically active components. Most of them have antioxidant activities (Hyo, 2012).

It is well known that under stressful conditions free radicals are over-produced. Oxidative stress occurs due to an imbalance between free radical formation and antioxidant defense capacity (Bahmani, 2014; Delfan, 2014; Nasri, 2014; Rafieian-Kopaei, 2013; Madihi, 2013). The result of this oxidative stress would be the induction of chronic hard curable diseases such as diabetes (Asgary, 2014; Rafieian-Kopaei, 2014; Rafieian-Kopaei, 2014; Nasri, 2014; Asgary, 2013), hypertension (Baradaran, 2014; Sadeghi, 2014), cardiovascular (Asgary, 2013; Madihi, 2013; Sarrafzadegan, 2013) cancer (Shirzad, 2009; 2011; 2013), cognitive diseases (Baradaran, 2012; Rabiei, 2013; Baradaran, 2012; Rabiei, 2014), and pain (Bahmani, 2014; Saki, 2014; Bahmani, 2014; Delfan, 2004) or exacerbation of some other diseases like infectious disorders (Asadbeygi, 2014; Bagheri, 2013; Bahmani, 2014; Bagheri, 2013; Bahmani, 2014; Amirmohammadi, 2014; Bahmani, 2014).

Synthetic antioxidants, in contrast to their natural counterparts, have been shown to produce side effects such as toxicity. Hence, it is important to prepare natural products with antioxidant activities that are able to prevent and treat free radical-associated diseases (Rahimian, 2014; Baradaran, 2014; Nasri, 2014; Nasri, 2014). A lot of plant constituents are known to have antioxidant activities (Nasri, 2014; Rafieian-Kopaei, 2014; Baradaran, 2014; Baradaran, 2014; Ghaed, 2014; Nasri, 2014; Bahmani, 2014; Nasri, 2013).

These plants have drawn much attraction because they have protective or curative properties against most of hard curable diseases such as cognitive deficit, memory impairment, cancer, and cardiovascular diseases (Taghikhani, 2014; Heidarian, 2013; Taghikhani, 2012; Roohafza, 2013; Parsaei, 2013; Asadi, 2013; Gharipour, 2013; Nasri, 2013; Gharipour, 2014; Gholami-Ahangaran, 2012; Bahmani, 2012; Gholami-Ahangaran, 2012; Bahmani, 2014; Bahmani, 2012; Bahmani, 2013; Eftekhari, 2013; Bahmani, 2012).

Hence, the therapeutic effects of licorice which has antioxidant activity on inflammation, malaria, cardiovascular disease, peptic ulcers, hepatitis C, and pulmonary, skin diseases, and so on might be, at least in part, attributed to its antioxidant activity. In this regard, there is ample evidence showing the ability of licorice is closely related to its antioxidant activities.

## REFERENCES

- Acharya SK, Dasarathy S, Tandon A, Joshi YK and Tandon BN, A preliminary open trial on interferon stimulator (SNMC) derived from *Glycyrrhiza glabra* in the treatment of subacute hepatic failure, Indian J. Med. Res., 98, 1993, 69 -74.
- Akhundzadeh S, Encyclopedia of Iranian Medicinal plants, Institute of Medicinal Plants, Jihad-e Daneshgahi, 2000, 213.
- Alan Teck WE, Yuan HM and Shi OE, Evaluation of surfactant assisted pressurized liquid extraction for the determination of glycyrrhizin and ephedrine in medicinal plants, Analytica Chimica Acta, 583, 2007, 289 - 95.
- Amirmohammadi M, Khajoenia S, Bahmani M, Rafieian-Kopaei M, Eftekhari Z, Qorbani M, *In vivo* evaluation of antiparasitic effects of *Artemisia abrotanum* and *Salvia officinalis* extracts on *Symphyla obvelata*, *Aspiculoris tetrapetra* and *Hymenolepis nana* parasites, Asian Pac J Trop Dis., 4(1), 2014, S250-S254.

Asadbeigi M, Mohammadi T, Rafieian-Kopaei M, Saki K, Bahmani M, Delfan B, Traditional effects of medicinal plants in the treatment of respiratory diseases and disorders: an ethnobotanical study in the Urmia, Asian Pac J Trop Med., 7(1), 2014, S364-S368.

Asadi SY, Parsaei P, Karimi M, Ezzati S, Zamiri A, Mohammadzadeh F, Rafieian-Kopaei M, Effect of green tea (*Camellia sinensis*) extract on healing process of surgical wounds in rat, Int J Surg., 11(4), 2013, 332-7.

Asadi-Samani M, Bahmani M, Rafieian-Kopaei M, The chemical composition, botanical characteristic and biological activities of *Borago officinalis*: a review, Asian Pac J Trop Med., 7(1), 2014, 22-28.

Asgary S, Kelishadi R, Rafieian-Kopaei M, Najafi S, Najafi M, Sahebkar A, Investigation of the lipid-modifying and antiinflammatory effects of *Cornus mas* L. supplementation on dyslipidemic children and adolescents, Pediatr Cardiol., 34(7), 2013.

Asgary S, Rafieian-Kopaei M, Shamsi F, Najafi S, Sahebkar A. Biochemical and histopathological study of the anti-hyperglycemic and anti-hyperlipidemic effects of cornelian cherry (*Cornus mas* L.) in alloxan-induced diabetic rats. J Complement Integr Med., 11(2), 2014, 63-9.

Asgary S, Sahebkar A, Afshani M, Keshvari M, Haghjooyjavanmard Sh, Mahmoud Rafieian-Kopaei M. Clinical evaluation of blood pressure lowering, endothelial function improving, hypolipidemic and anti-inflammatory effects of pomegranate juice in hypertensive subjects. Phytother. Res., 2013.

Bagheri N, Rahimian Gh, Salimzadeh L, Azadegan F, Rafieian-Kopaei M, Taghikhani A, Shirzad H, Association of the Virulence factors of *Helicobacter pylori* and Gastric Mucosal Interleukin-17/23 mRNA Expression in dyspeptic patients. EXCLI J., 12, 2013, 5-14.

Bagheri N, Taghikhani A, Rahimian G, Salimzadeh L, Azadegan Dehkordi F, Zandi F, Chaleshtori MH, Rafieian-Kopaei M, Shirzad H. Association between virulence factors of *helicobacter pylori* and gastric mucosal interleukin-18 mRNA expression in dyspeptic patients. Microb Pathog., 2013.

Bahmani M, Farkhondeh T and Sadighara P. The anti-parasitic effects of *Nicotina tabacum* on leeches, Comparative Clinical Pathology, 21(3), 2012, 357-359.

Bahmani M, Forouzan Sh, Rafieian-Kopaei M, Eftekhari Z, Evaluating the Anti-Leech Effects of Methanolic Extracts of *Peganum harmala* L. and *Olea europaea* L. on immature worm *Limnatis nilotica*, Asian Pac J Trop Dis., 2012, 1-6.

Bahmani M, Karamati SA, Banihabib EK, Saki K, Comparison of effect of nicotine and levamisole and ivermectin on mortality of leech, Asian Pac J Trop Dis., 4(1), 2014, 477-480.

Bahmani M, Karamati SA, Hassanzadazar H, Forouzan SH, Rafieian-Kopaei M, Kazemi-Ghoshchi B, Asadzadeh J, Kheiri AG, Ehsan Bahmani E, Ethnobotanic study of medicinal plants in Urmia city: identification and traditional using of antiparasites plants, Asian Pac J Trop Dis., 4(2), 2014, 906-910.

Bahmani M, Rafieian M, Baradaran A, Rafieian S, Rafieian-kopaei M, Nephrotoxicity and hepatotoxicity evaluation of *Crocus sativus* stigmas in neonates of nursing mice, J Nephrothol., 3(2), 2014, 81-85.

Bahmani M, Rafieian-Kopaei M, Hassanzadazar H, Saki K, Karamati SA, Delfan B, A review on most important herbal and synthetic antihelmintic drugs, Asian Pac J Trop Med., 7(1), 2014, 29-33.

Bahmani M, Rafieian-Kopaei M, Jeloudari M, Eftekhari Z, Delfan B, Zargaran A, Forouzan SH, A review of the health effects and uses of drugs of plant licorice (*Glycyrrhiza glabra* L.) in Iran, Asian Pac J Trop Dis., 4(2), 2014, 847-849.

Bahmani M, Saki K, Gholami-Ahangaran M, Parsaei P, Mohsenzadegan A, Zia-Jahromi N, Evaluating the Anti-Leech Activity of Methanolic Extract of *Matricaria chamomilla* L. Comparing with Ivermectin, Mebendasole, Praziquantel, Rafoxanide, Febantel and Albendazole, Mid East J Sci Res., 12(2), 2012, 260-263.

Bahmani M, Saki k, Rafieian-Kopaei M, Karamati SA, Eftekhari Z, Jelodari M, The most common herbal medicines affecting *Sarcocystis* branches: a review study, Asian Pac.J.Trop.Biomed., 4(12), 2014, 930-937.

Bahmani M, Shirzad HA, Majlesi M, Shahinfard N, Rafieian-Kopaei M, A review study on analgesic applications of Iranian medicinal plants, Asian Pac J Trop Med., 7(1), 2014, 43-53.

Bahmani M, Vakili-Saatloo N, Gholami-Ahangaran M, Karamati SA, Khalil-Banihabib E, Hajjigholizadeh Gh, A comparison study on the anti-leech effects of onion (*Allium cepa* L) and ginger (*Zingiber officinale*) with levamisole and triclabendazole, J HerbMed Pharmacol., 2(1), 2013, 1-3.

- Bahmani M, Zargaran A, Rafieian-Kopaei M, Saki M, Ethnobotanical study of medicinal plants used in the management of diabetes mellitus in the Urmia, Northwest Iran. *Asian Pac J Trop Med.*, 7(1), 2014, 348-354.
- Bahmani M, Zargaran A, Rafieian-Kopaei M, Identification of medicinal plants of Urmia for treatment of gastrointestinal disorders, *Rev Bras Farmacogn.*, 24, 2014, 468.
- Bahmani, M., Rafieian-Kopaei, M, Medicinal plants and secondary metabolites for leech control, *Asian Pac J Trop Dis.*, 2014; 4(4), 2014, 315-316.
- Baradaran A, Nasri H, Nematbakhsh M, Rafieian-Kopaei M, Antioxidant activity and preventive effect of aqueous leaf extract of Aloe Vera on gentamicin-induced nephrotoxicity in male Wistar rats, *Clin Ter.*, 165(1), 2014, 7-11.
- Baradaran A, Nasri H, Rafieian-Kopaei M, Comment on: Anti-oxidative stress activity of *Stachys lavandulifolia* aqueous extract in humans, *Cell J.*, 15(3), 2013, 272-3.
- Baradaran A, Nasri H, Rafieian-Kopaei M, Oxidative stress and hypertension: Possibility of hypertension therapy with antioxidants, *J Res Med Sci.*, 19(4), 2014, 358-67.
- Baradaran A, Rabiei Z, Rafieian M, Shirzad H, A review study on medicinal plants affecting amnesia through cholinergic system, *Journal of HerbMed Pharmacology*, 1(1), 2012, 3-9.
- Baradaran A, Madihi Y, Merrikhi A, Rafieian-Kopaei M, Nematbakhsh M, Asgari A, Khosravi Z, Haghghian F, Nasri H, Nephrotoxicity of hydroalcoholic extract of *Teucrium polium* in Wistar rats, *Pak J Med Sci.*, 29(1), 2013, 329-333.
- Blumenthal M, Goldberg A and Brinckmann J, *Herbal Medicine, Expanded Commission E Monographs*, 1st ed., Integrative Medicine Communications, USA, 2000, 233 - 5.
- British Pharmacopeia, Stationery Office, London, 2, 2004, 1175 – 6.
- Chandler F, *Herbs Everyday Reference for Health professionals*, Canadian Pharmacists Association and the Canadian Medicinal Association, 2000.
- Colalto C, Herbal in traction on absorption of drugs: Mechanisms of action and clinical risk assessment, *Pharmacol. Res.*, 62, 2010, 207 - 27.
- Csuk R, Schwarz S and Kluge R, Synthesis and biological activity of some anti tumora active derivatives from glycyrrhetic acid, *European Journal of Medicinal Chem.*, 45, 2010, 5718 - 23.
- Delfan B, Bahmani M, Eftekhari Z, Jelodari M, Saki K, Mohammadi T, Effective herbs on the wound and skin disorders: a ethnobotanical study in Lorestan province, west of Iran, *Asian Pac J Trop Dis.*, 4(2), 2014, 938-942.
- Delfan B, Bahmani M, Hassanzadazar H, Saki K, Rafieian-Kopaei M, Identification of medicinal plants affecting on headaches and migraines in Lorestan Province, West of Iran, *Asian Pac J Trop Med.*, 7(1), 2014, 376-379.
- Delfan B, Bahmani M, Rafieian-Kopaei M, Delfan M, Saki K, A review study on ethnobotanical study of medicinal plants used in relief of toothache in Lorestan Province, Iran, *Asian Pac J Trop Dis.*, 4(2), 2014, 879-884.
- Douglas JA, Douglas MH, Lauren DR, Martin RJ, Deo B, Follett JM and Jensen DJ, Effect of plant density and depth of harvest on the production and quality of licorice (*Glycyrrhiza glabra*) root harvested over 3 years, *NZJC*, 32, 2004, 363 - 73.
- Eftekhari Z, Bahmani M, Mohsenzadegan A, Gholami-Ahangaran M, Abbasi J, Alighazi N, Evaluating the anti-leech (*Limnatis nilotica*) activity of methanolic extract of *Allium sativum* L. compared with levamisole and metronidazole, *Comp Clin Path.*, 21, 2012, 1219-1222.
- Farnsworth NR, Fong HHS, Malady GB, Gummi Gugguli, WHO monographs on selected medicinal plants, Volume III, WHO Traditional Medicinal programme, Geneva, Switzerland, 2004.
- Fenwick GK, Lutomski J and Nieman C, Licorice, *Glycyrrhiza glabra* L. composition, uses and analysis, *Food Chem.*, 38, 1990, 119 - 43.
- Fiore C, Eisenhut M, Krausse R, Ragazzi E, Pellati D, Armanini D and Bielenberg J, Antiurial effects of *Glycyrrhiza* species, *Phytotherapy Res.*, 22, 2008, 141 - 8.
- Fukai T, Marumoa A, Kaitou K, Kanda T, Terada S and Nomura T, Anti-Helicobacter pylori flavonoids from licorice extract, *Life Sciences*, 71(12), 2002, 1449 – 63.

Gazzani G, Daglia M, Papetti A, Food components with anticaries activity, *Current Opinion in Biotechnology*, 23(2), 2012, 153–9.

Ghaed F, Rafieian-Kopaei M, Nematbakhsh M, Baradaran A, Nasri H, Ameliorative effects of metformin on renal histologic and biochemical alterations of gentamicin-induced renal toxicity in Wistar rats, *J Res Med Sci.*, 17(7), 2012, 621-625.

Ghahraman A, *Basic Botany: Anatomy and Morphology*, Vol.1, University of Tehran Press, 1999, 539.

Ghanbari A, Rahimiyan Mashhadi H, Nasiri mahalati M and Rastgo M, Ecophysiological aspects of licorice germination response to temperature, *Journal of Agricultural Res.*, 3 (2), 2005, 263 - 75.

Gharipour M, Ramezani MA, Sadeghi M, Khosravi A, Masjedi M, Khosravi-Boroujeni H, Sex based levels of C-reactive protein and white blood cell count in subjects with metabolic syndrome: Isfahan Healthy Heart Program, *J Res Med Sci.*, 18, 2013, 467-72.

Gholami-Ahangaran M, Bahmani M, Zia-Jahrom N, *In vitro* antileech effects of *Vitis vinifera* L., niclosamide and ivermectin on mature and immature forms of leech *Limnatis nilotica*, *Glob Vet*, 8, 2012, 229-232.

Gholami-Ahangaran M, Bahmani M, Zia-Jahromi N, Comparative and evaluation of anti-leech (*Limnatis Nilotica*) effect of Olive (*Olea Europaea* L.) with levamisole and tiabendazole, *Asian Pac J Trop Dis.*, 2(1), 2012, 101-103.

Hayashi H, Hiraoka N, Ikeshiro Y, Yammamoto H and Yoshikawa T, Seasonal variation of glycyrrhizin and isoliquiritigenin glycosides in the root of *Glycyrrhiza glabra* L., *Biological and Pharmaceutical Bulletin*, 21(9), 1998, 987 - 9.

Heidarian E, Rafieian-Kopaei M, Protective effect of artichoke (*Cynara scolymus*) leaf extract against lead toxicity in rat, *Pharm Biol.*, 51(9), 2013, 1104-9.

Hyo- Jung K, Ji-Yeon S, Hwa-Jin S, Soon- Sung, Jong-Sang K, Antioxidant activities of licorice-derived phenyl flavonoids, *Nutr Res Pract.*, 6(6), 2012, 491-498.

Jiang Y, Lu TH and Chen F, Preparative purification of glycyrrhizin extracted from the root of licorice using high-speed counter-current chromatography, *J. Chromatogr.*, 1033, 2004, 183-6.

Karamati SA, Hassanzadazar H, Bahmani M, Rafieian-Kopaei M, Herbal and chemical drugs effective on malaria, *Asian Pac J Trop Dis.*, 4(2), 2014, 599-601.

Khosravi-Boroujeni H, Mohammadifard N, Sarrafzadegan N, Sajjadi F, Maghroun M, Khosravi A, Alikhasi H, Rafieian M, Azadbakht L, Potato consumption and cardiovascular disease risk factors among Iranian population, *Int J Food Sci Nutr.*, 63(8), 2012, 913-20.

Khosravi-Boroujeni H, Sarrafzadegan N, Mohammadifard N, Sajjadi F, Maghroun M, Asgari S, Rafieian-Kopaei M, Azadbakht L, White rice consumption and CVD risk factors among Iranian population, *J Health Popul Nutr.*, 31(2), 2013, 252-61.

Li w, Asada Y and Yoshikawa T, Flavonoid constituents from *Glycyrrhiza glabra* hairy root cultures, *Phytochem*, 55, 2000, 447 - 56.

Li YJ, Chen J, Li Y, Li Q, Zheng YF, Fu Y and Li P, Screening and characterization of natural antioxidants in four *Glycyrrhiza* species by liquid chromatography coupled with electrospray ionization quadrupole time of flight tandem mass spectrometry, *J. Chromatogr A*, 1218 (45), 2011, 8181 - 91.

Madihi Y, Merrikhi A, Baradaran A, Rafieian-kopaei M, Shahinfard N, Ansari R, Shirzad H, Mesripour A, Impact of sumac on postprandial high-fat oxidative stress, *Pak J Med Sci.*, 29 (1), 2013, 340-345.

Madihi Y, Merrikhi A, Baradaran A, Ghobadi S, Shahinfard N, Ansari R, Karimi A, Mesripour A, Rafieian-Kopaei M, Bioactive components and the effect of hydroalcoholic extract of *Vaccinium myrtillus* on postprandial atherosclerosis risk factors in rabbits, *Pak J Med Sci.*, 29(1), 2013, 384-389.

Maha M, Gazial A and Nermeen M, Effect of Glabridin on the Structure of Ileum and Pancreas in Diabetic Rats: A histological, Immunohistochemical and Ultrastructural Study, *Nature and Sci.*, 10(3), 2012, 78 - 90.

Merriam-Webster's Medical Dictionary, Merriam-Webster, Inc. 2007.

Messier C, Epifano F, Genovese S, Grenier D, Licorice and its potential beneficial effects in common oro-dental diseases, *Oral Diseases*, 18 (1), 2012, 32–9.



Mirhaidar H, Licorice, Herbal plants used in the treatment of diseases and education, Office of Islamic Culture Publication, 3, 1994, 12 -16.

Mirhosseini M, Baradaran A, Rafieian-Kopaei M, Anethum graveolens and hyperlipidemia: A randomized clinical trial, J Res Med Sci., 19, 2014, 758-61.

Mojtahedi M and Lesani M, Life of Green plants, University of Tehran Press, 2015, 587.

Montoro P, Maldini M, Russo M, Postorino S, Piacente S and Pizza C, Metabolic profiling of roots of liquorice (*Glycyrrhiza glabra*) from different geographical areas by ESI/MS/MS and determination of major metabolites by LC-ESI/MS and LC-ESI/MS/MS, J.Pharm.Biomed Anal., 54, 2011, 535 - 45.

Nasri H, Baradaran A, Ardalan MR, Mardani S, Momeni A, Rafieian-Kopaei M, Bright renoprotective properties of metformin: beyond blood glucose regulatory effects, Iran J Kidney Dis., 7(6), 2013, 423-8.

Nasri H, Nematbakhsh M, Rafieian-Kopaei M, Ethanolic extract of garlic for attenuation of gentamicin-induced nephrotoxicity in Wistar rats, Iran J Kidney Dis., 7(5), 2013, 376-82.

Nasri H, Rafieian-Kopaei M, Medicinal plants and antioxidants: Why they are not always beneficial?, Iranian Journal of Public Health, 43(2), 2014, 255-257.

Nasri H, Rafieian-Kopaei M, Medicinal plants and new concerns in statin consumption, Iranian J Publ Health, 42(9), 2013, 1071-1072.

Nasri H, Rafieian-Kopaei M, Protective effects of herbal antioxidants on diabetic kidney disease, J Res Med Sci., 19(1), 2014, 82-3.

Nasri H, Tavakoli M, Ahmadi A, Baradaran A, Nematbakhsh M, Rafieian-Kopaei M, Ameliorative effect of melatonin against contrast media induced renal tubular cell injury, Pak J Med Sci., 30(2), 2014, 261-5.

Nasri H, Rafieian-Kopaei M, Tubular kidney protection by antioxidants, Iranian J Publ Health, 42(10), 2013, 1194-1196.

Nezamabadi H, Rahimiyan Mashhadi H, Zand A and Alizadeh H, Ecophysiological aspects of Licorice rhizome, Plant Diseases and Pests, 74 (2), 2006, 45 – 62.

Omar-Hesham R, Komarova I, El-Ghonemi MA, Fathy-Rashad R, Abdelmalak HD, Yerramadha MR, How much is too much? in Licorice abuse: time to send a warning message from Therapeutic Advances in Endocrinology and Metabolism, <http://www.ncbi.nlm.nih.gov>, SAGE Publications, Retrieved 13 January 2015.

Omidbeigi R, Processing and production of medicinal plants, Razavi Publications, Mashhad, 3, 2006, 397.

Ong ES and Len SM, Pressurized hot water extraction of berberine and baicalin and glycyrrhizin in medicinal plants, Analytica Chimica Acta, 482, 2003, 81 - 9.

Parsaei P, Karimi M, Asadi SY, Rafieian-Kopaei M, Bioactive components and preventive effect of green tea (*Camellia sinensis*) extract on postlaparotomy intra-abdominal adhesion in rats, Int J Surg., 2013.

Rabiei Z, Hojjati M, Rafieian-Kopaei M, Alibabaei Z, Effect of *Cyperus rotundus* tubers ethanolic extract on learning and memory in animal model of Alzheimer, Biomedicine & Aging Pathology, 3(4), 2013, 185-91.

Rabiei Z, Rafieian-Kopaei M, Mokhtari S, Alibabaei Z, Shahrani M, The effect of pretreatment with different doses of *Lavandula officinalis* ethanolic extract on memory, learning and nociception, Biomedicine & Aging Pathology, 4(1), 2014, 71-6.

Rafieian-Kopaei M, Baradaran A, Rafieian M, Oxidative stress and the paradoxical effects of antioxidants, J Res Med Sci., 18(7), 2013, 628.

Rafieian-Kopaei M, Behradmanesh S, Kheiri S, Nasri H, Association of serum uric Acid with level of blood pressure in type 2 diabetic patients, Iran J Kidney Dis., 8(2), 2014, 152-4.

Rafieian-Kopaei M, Nasri H, Re: Erythropoietin ameliorates oxidative stress and tissue injury following renal ischemia/reperfusion in rat kidney and lung, Med Princ Pract., 23(1), 2014, 95.

Rafieian-Kopaei M, Nasri H, The Ameliorative Effect of *Zingiber officinale* in Diabetic Nephropathy, Iran Red Crescent Med J., 16(5), 2014, e11324.

Rafieian-Kopaei M, Setorki M, Doudi M, Baradaran A, Nasri H, Atherosclerosis: Process, Indicators, Risk Factors and New Hopes, Int J Prev Med., 5, 2014, 927-46.

- Rafieian-Kopaei M, Shahinfard N, Rouhi-Boroujeni H, Gharipour M, Darvishzadeh-Boroujeni P, Effects of *Ferulago angulata* extract on serum lipids and lipid peroxidation, *Evid Based Complement Alternat Med.*, 2014.
- Rahimian G, Sanei MH, Shirzad H, Azadegan-Dehkordi F, Taghikhani A, Salimzadeh L, Hashemzadeh-Chaleshtori M, Rafieian-Kopaei M, Bagheri N, Virulence factors of *Helicobacter pylori* vacA increase markedly gastric mucosal TGF- $\beta$ 1 mRNA expression in gastritis patients, *Microb Pathog.*, 2014, 67-68.
- Rahnama S, Rabiei Z, Alibabaei Z, Mokhtari S, Rafieian-kopaei M, Deris F, Anti-amnesic activity of *Citrus aurantium* flowers extract against scopolamine-induced memory impairments in rats, *Neurological Sciences*, 2014, 1-8.
- Rengel Z and Graham RD, Uptake of Zinc from chelate- buffered nutrient by wheat genotype differing Zn deficiency, *JXB*, 47, 1996, 217 - 26.
- Roohafza H, Sarrafzadegan N, Sadeghi M, Rafieian-Kopaei M, Sajjadi F, Khosravi-Boroujeni H, The association between stress levels and food consumption among Iranian population, *Arch Iran Med.*, 16(3), 2013, 145-8.
- Sabbioni C, Ferranti A, Bugamelli F, Cantelli Forti G and Augusta Raggi M, Simultaneous HPLC analysis, with isocratic elution, of glycyrrhizin and glycyrrhetic acid in licorice root an confectionary products, *Phytochemical Analysis*, 17, 2005, 25 - 31.
- Sadeghi M, Khosravi-Boroujeni H, Sarrafzadegan N, Asgary S, Roohafza H, Gharipour M, Sajjadi F, Khalesi S, Rafieian-Kopaei M, Cheese consumption in relation to cardiovascular risk factors among Iranian adults- IHHP Study, *Nutr Res Pract.*, 8(3), 2014, 336-41.
- Saki K, Bahmani M, Rafieian-Kopaei M, Hassanzadazar H, Dehghan K, Bahmani F, Asadzadeh J, The most common native medicinal plants used for psychiatric and neurological disorders in Urmia city, northwest of Iran, *Asian Pac J Trop Dis.*, 4(2), 2014, 895-901.
- Saki K, Bahmani M, Rafieian-Kopaei M, The effect of most important medicinal plants on two important psychiatric disorders (anxiety and depression)-a review, *Asian Pac J Trop Med.*, 7(1), 2014, 34-42.
- Sarrafzadegan N, Khosravi-Boroujeni H, Esmailzadeh A, Sadeghi M, Rafieian-Kopaei, M., Asgary S, The association between hypertriglyceridemic waist phenotype, menopause, and cardiovascular risk factors, *Arch Iran Med.*, 16(3), 2013, 161-166.
- Shi OE and Mei LS, Pressurized hot water extraction of berberin, baicalein and glycyrrhizin in medicinal plants, *Analytica Chimica Acta*, 482, 2003, 81 - 9.
- Shirzad H, Kiani M, Shirzad M, Impacts of tomato extract on the mice fibrosarcoma cells, *J HerbMed Pharmacol.*, 2(1), 2013, 13-16.
- Shirzad H, Shahrani M, Rafieian-Kopaei M, Comparison of morphine and tramadol effects on phagocytic activity of mice peritoneal phagocytes *in vivo*, *Int Immunopharmacol.*, 9(7-8), 2009, 968-70.
- Shirzad H, Taji F, Rafieian-Kopaei M, Correlation between antioxidant activity of garlic extracts and WEHI-164 fibrosarcoma tumor growth in BALB/c mice, *J Med Food*, 14(9), 2011, 969-74.
- Somjen D, Katzburg S, Vaya J, Kaye AM, Hendel D, Posner G.H, Tamir S, Estrogenic activity of glabridin and glabrene from licorice roots on human osteoblasts and prepubertal rat skeletal tissues, *The Journal of Steroid Biochemistry and Molecular Biology*, 91 (4-5): 241-246.
- Taghikhani A, Afrough H, Ansari-Samani R, Shahinfard N, Rafieian-Kopaei M, Assessing the toxic effects of hydroalcoholic extract of *Stachys lavandulifolia* Vahl on rat's liver, *Bratisl Lek Listy*, 115(3), 2014, 121-4.
- Taghikhani M, Nasri H, Asgari A, Afrough H, Namjoo AR, Ansari-Samani R, Shahinfard N, Rafieian-kopaei M, The renal toxicity of hydroalcoholic extract of *Stachys lavandulifolia* Vahl in Wistar rats, *Life Sci J.*, 9(4), 2012, 3025-31.
- Tamir S, Eizenberg M, Somjen D, Izrael S, Vaya J, Estrogen-like activity of glabrene and other constituents isolated from licorice root, *The Journal of steroid biochemistry and molecular biology*, 78 (3), 2001, 291-298.
- USP 28/ NF 23, The united States Pharmacopeial Convention, Toronto, 3, 2005, 2109 – 10.
- Waliullahpoor R, Rashed Mohassel H M, Effect of planting depth and size of the rhizome on liquorice growth, *Journal of Agriculture and Natural Resources*, 12, 2005, 1-12.