




Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Disease

journal homepage: www.elsevier.com/locate/apjtd

Document heading

doi:10.1016/S2222-1808(14)60580-6

© 2014 by the Asian Pacific Journal of Tropical Disease. All rights reserved.

Medicinal plants and secondary metabolites for leech control

Mahmoud Bahmani^{1,2*}, Mahmoud Rafeian-Kopaei³¹Razi Herbal Medicine Research Center, Lorestan University of Medical Sciences, Khorram Abad, Iran²Food and Beverages Safety Research Center, Urmia University of Medical Sciences, Urmia, Iran³Medical Plants Research Center, Shahrekord University of Medical sciences, Shahrekord, Iran

Dear editor,

Leech is a parasitic animal that belongs to the phylum Annelida. Some species of Leeches are aquatic and live in freshwater rivers and lakes, marshes and paddy fields and some other species live in terrestrial and marine environments. Leeches are necrophagous, predatory or parasitic animals and a group of them Hematophagy habit. The blood sucking leech attaches to the skin and other internal organs of the body like mouth, throat, trachea and vagina by cupping and a painless bit and holes its skin or phlegm and secretes an anticoagulant enzyme into the hosts' blood streams. Each leech secretes 5–15 mL hirudin in one meal and sucks the blood[1–3]. They, as a harmful parasite can cause some side effects like severe anemia, short-term hemorrhage, hypersensitivity and anaphylactic reactions[1,4]. Most physicians who have practiced in remote and rural areas of Iran have complete familiarity with the leech as a foreign body in nose, nasopharynx, pharynx and larynx and also the respiration system[5]. Leech is one of the factors of expressing zoonotic diseases[6]. Leeches are parasite of the human, livestock, turtles, frogs, snails and earth worms and feed from them. The parasitic potential of the leech is high to the extent that even sluedges in bottom of the pool are not safe from their attacks[1,3,4]. No standard drug has been registered in the pharmacopeia of different nations in order to treat effects of leech infestation and treatment are supportive and traditional including the use of hypertonic saline, lidocaine and vinegar that are used to remove and the leech from the bit position[7–10]. For example, wearing socks when feet are exposed to the risk, putting dried remains of the bath soap on the bite site, putting tobacco leaves between fingers of the toe, placing some salt or bicarbonate, citrus extract, eucalyptus oil, washing place of the bit with

water and soap, using a bag of ice water and cooling place to decrease pain and inflammation, using antihistamine to decrease itching and inflammation and even using calcium hydroxide causes the tissue irritation and damage. These are traditional treatments used by people in different culture of the world in cases of infection with the leech and are not effective treatments[11,12]. Since the ancient time using pharmaceutical plants to treat parasitic diseases has been paid attention. On one hand due to appearing adverse and undesirable effects of synthetic compounds and their incompatibility with human nature and on other hand due to high cost and drugs and the inability of many third world countries to purchase such drugs a special attention has been paid to produce drug from pharmaceutical plants that their safe therapeutic effects and inexpensive cost of them have been proved[13,14].

The leech is inherently aggressive and causes much infection. There are several reports about different animal infections including cow, sheep, dandy, hen, dog, goat and lamb with the leech[15–17]. By the end of the year 2009 no scientific study was conducted about the effect of pharmaceutical plants on leech. It has stated in the resources of the Iran traditional medicine that the tobacco plant has anti-leech effect. In the traditional medicine decoction of the tobacco plant is used to repel leech[18]. Also the *Satureja khuzistanica* Jamzed plant has been defined as the anti-leech plant[18]. In recent years authors have examined the effect of various extracts of many local pharmaceutical plants on the *Limnatis nilotica* leech. It has been determined that the tobacco plant has the anti-leech and strong disinfection effect on the leech[6,19]. The nicotine substance, one of the effective tobacco substances, has very strong effect on the leech death[20,21]. The *Allium sativum* L. extract has suitable cytotoxicity effect on the leech[22,23].

*Corresponding author: Dr. Mahmoud Bahmani, Razi Herbal Medicine Research Center, Lorestan University of Medical Sciences, Khorram Abad, Iran.

Tel: 0989186157084

E-mail: mahmood.bahmani@gmail.com

Article history:

Received 11 Jun 2014

Received in revised form 19 Jun, 2nd revised form 25 Jun, 3rd revised form 5 Jul 2014

Accepted 23 Jul 2014

Available online 28 Aug 2014

Also the olive plant extract has a good effect on the leech death and paralysis^[24]. The *Zingiber officinale* is of plants that has the disinfection and anti leech effect^[25,26]. The *Vitis vinifera* has been determined as a plant with the antiparasitic effect on leech^[27]. Afsintin (*Artemisia* spp.) causes the leech death^[28]. Also *Artemisia kermanensis* has the strong effect on the leech death^[28]. *Matricaria chamomilla* is an anti leech plant^[29]. It seems that anti-parasitic plants have the suitable effect due to having secondary active compounds. With regard to importance of Leech infestation, it seems necessary to conduct applied and pharmacologic researches related to this subject in order to produce natural and effective drugs.

Conflict of interest statement

We declare that we have no conflict of interest.

References

- [1] Haycox CL, Odland PB, Clotrea MD, Raugi CJ. Indications and complications of medicinal leech therapy. *J Am Acad Dermatol* 1995; **33**: 1053–1055.
- [2] Vera Blu A, Torres M. [Leeches, today and yesterday present and past]. *Rev Chilena Infectol* 2005; **22**: 32–37. Spanish.
- [3] Will RB. Hirudins and the role thrombin: lessons from leeches. *Trends Pharmacol Sci* 1988; **9**: 425–427.
- [4] Reza SN, Shabbir SM, Anwar-ul-Haq. Leech infestation and its association with water drinking habits. *J Coll Physicians Surg Pak* 2006; **16**: 175–178.
- [5] Mohammad Y, Rostam M, Dubaybo BA. Laryngeal hirudiniasis: an unusual cause of airway obstruction and hemoptysis. *Pediatr Pulmonol* 2002; **33**: 224–226.
- [6] Bahmani M, Avijgan M, Hosseini SR, Qorbani M. Evaluating the anti-*Limnatis nilotica* effects of tobacco methanol extract compared with succinyl choline and some other anti-parasitic drugs. *Shahrekord J Med Sci* 2010; **12**(3): 53–59.
- [7] White GB. Leeches and leech infestation. In: Cook GC, editor. *Manson's tropical diseases*. London: Saunders; 2002, p. 1599–1600.
- [8] Litch JA, Bishop RA. Saturated aqueous sodium chloride solution for the removal of leeches. *Trop Doct* 2000; **30**: 102.
- [9] Leeches. *Lancet* 1825; **5**: 427.
- [10] Kruger C, Malleyeck I, Olsen OH. Aquatic leech infestation: a rare cause of severe anaemia in an adolescent Tanzanian girl. *Eur J Pediatr* 2004; **163**: 297–299.
- [11] Madill J, Hovingh P. Freshwater leech (Annelida: Hirudinida) distribution in the Canadian Province of Newfoundland and Labrador and adjacent regions: check-list, new records, new pigmentation forms, and *Pleistocene refugia*. *Zootaxa* 2007; **1657**: 1–21.
- [12] Schenkova J, Sychra J, Kubova B. The freshwater leeches (Clitellata: Hirudinida) of the Czech Republic—preference of different types of water bodies. In: 4th International Conference Hirudinea: biology, taxonomy, faunistics. 2011.
- [13] Nasri H, Shirzad H. Toxicity and safety of medicinal plants. *J HerbMed Pharmacol* 2013; **2**(2): 21–22.
- [14] Rafieian-Kopaei M. Medicinal plants and the human needs. *J HerbMed Pharmacol* 2012; **1**(1): 1–2.
- [15] Bahmani M, Golshahi H, Ghotbian F, Bahmani F. Internal hirudiniasis in a hen (*Gallus gallusdomesticus*)—the first report in literature. *Asian Pac J Trop Dis* 2013; **3**: 232–234.
- [16] Bahmani M, Eftekhari Z, Mohsenzadeghan A, Ghotbian F, Alighazi N. Leech (*Limnatis nilotica*) causing respiratory distress in a pregnant cow in Ilam province in Iran. *Comp Clin Pathol* 2012; **21**: 501–503.
- [17] Bahmani M, Rasouli R, Parsaei P, Bahnihabib E, Saki K, Ghotbian F. *Limnatis nilotica* infestation in ram and kid in Dehloran city, Ilam province, west of Iran. *Asian Pac J Trop Dis* 2013; **3**(2): 155–157.
- [18] Bahmani M, Eftekhari Z. An ethnoveterinary study of medicinal plants in treatment of diseases and syndromes of herd dog in southern regions of Ilam province, Iran. *Comp Clin Pathol* 2012; **22**: 403–407.
- [19] Bahmani M, Farkhondeh T, Sadighara P. The anti-parasitic effects of *Nicotina tabacum* on leeches. *Comp Clin Pathol* 2012; **21**(3): 357–359.
- [20] Bahmani M, Karamati SA, Banihabib E, Saki K. Comparison of effect of nicotine and levamisole and ivermectin on mortality of leech. *Asian Pac J Trop Dis* 2014; **4**(Suppl 1): S477–S480.
- [21] Bahmani MK, Banihabib E. Comparative assessment of the anti-annelida (*Limnatis nilotica*) activity of nicotine with niclosamide. *Global Veterinaria* 2013; **10**(2): 153–157.
- [22] Eftekhari Z, Bahmani M, Mohsenzadegan A, Gholami-Ahangaran M, Abbasi J, Alighazi N. Evaluating the anti-leech (*Limnatis nilotica*) activity of methanolic extract of *Allium sativum* L. compared with levamisole and metronidazole. *Comp Clin Pathol* 2012; **21**: 1219–1222.
- [23] Bahmani M, Abbasi J, Mohsenzadegan A, Sadeghian S, Gholami-Ahangaran M. *Allium sativum* L. the anti-immature leech (*Limnatis nilotica*) activity compared to niclosamide. *Comp Clin Pathol* 2011; doi: 10.1007/s00580-011-1380-7.
- [24] Gholami-Ahangaran M, Bahmani M, Zia-Jahromi N. Comparative and evaluation of anti-leech (*Limnatis Nilotica*) effect of olive (*Olea Europaea* L.) with levamisole and tiabendazole. *Asian Pac J Trop Dis* 2012; **2**(Suppl 1): S101–S103.
- [25] Bahmani M, Golshahi H, Mohsenzadegan A, Gholami-Ahangaran M, Ghasemi E. Comparative assessment of the anti-*Limnatis nilotica* activities of *Zingiber officinale* methanolic extract with levamisole. *Comp Clin Pathol* 2013; **22**(4): 667–670.
- [26] Forouzan S, Bahmani M, Parsaei P, Mohsenzadegan A, Gholami-Ahangaran M, Sadeghi E, et al. Anti-parasitic activities of *Zingiber officinale* methanolic extract on *Limnatis nilotica*. *Global Veterinaria* 2012; **9**(2): 144–148.
- [27] Gholami-Ahangaran M, Bahmani M, Zia-Jahromi N. *In vitro* antileech effects of *Vitis vinifera* L., niclosamide and ivermectin on mature and immature forms of leech *Limnatis nilotica*. *Global Veterinaria* 2012; **8**: 229–232.
- [28] Bahmani M, Avijgan M, Gholami-Ahangaran M, Rafieian M. The comparison of anti *Limnatis nilotica* effects of albendazole and some of the Iranian medicinal plants. *Iran South Med J* 2012; **15**(1): 25–34.
- [29] Bahmani M, Saki K, Gholami-Ahangaran M, Parsaei P, Mohsenzadegan A, Zia-Jahromi N. Evaluating the anti-leech activity of methanolic extract of *Matricaria chamomilla* L. comparing with ivermectin, mebendasole, praziquantel, rafoxanide, febantel and albendasole. *Middle-East J Sci Res* 2012; **12**(2): 260–263.