

# Protective effects of *Cornus mas* fruit extract on carbon tetrachloride induced nephrotoxicity in rats

M. Es.Haghi, G. Dehghan<sup>1</sup>, N. Banihabib<sup>2</sup>, S. Zare<sup>3</sup>, P. Mikaili<sup>4</sup>, F. Panahi<sup>5</sup>

Liver and Gastrointestinal Diseases Research Center, <sup>1</sup>Department of Biology, Faculty of Natural Sciences, University of Tabriz, <sup>2</sup>Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, <sup>3</sup>Department of Biology, Faculty of Science, University of Urmia, <sup>4</sup>Department of Pharmacology, School of Medicine, Urmia University of Medical Sciences, Urmia, <sup>5</sup>Neurosciences Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

## ABSTRACT

Oxidative damage is implicated in the pathogenesis of kidney injury. *Cornus mas* is used for in renal ailments traditionally in Iran. The present study was aimed to investigate the antioxidant activity of *C. mas* fruit extract (CMFE) on carbon tetrachloride (CCl<sub>4</sub>) treated oxidative stress in Wistar albino rats. Forty two male albino rats were divided into seven groups. Group I served as a sham; Group II served as a normal control; Group III served as a toxic control, with CCl<sub>4</sub> (1 ml/kg body weight; 80% in olive oil); Groups IV and V received CMFE at doses of 300 and 700 mg/kg before CCl<sub>4</sub> injection; Groups VI and VII received extract at same doses orally at 2, 6, 12, 24 and 48 h after CCl<sub>4</sub> intoxication. CCl<sub>4</sub> injection produced a significant rise in serum markers of oxidative stress and lipid peroxidation product malondialdehyde along with the reduction of antioxidant enzymes such as superoxide dismuta, catalase and glutathion peroxidase. Serum creatinine, urea and uric acid concentrations were increased whereas level of protein and albumin were reduced. Treatment of rats with different doses of fruit extract (300 and 700 mg/kg) significantly ( $P < 0.05$ ) ameliorated the alterations induced with CCl<sub>4</sub> in lipid peroxidation, antioxidant defenses, biochemical and renal lesions. Based on these results, we conclude that CMFE protects kidney from oxidative stress induced by CCl<sub>4</sub>.

**Key words:** Carbon tetrachloride, *Cornus mas*, lipid paroxidation, nephrotoxicity, oxidative stress

## Introduction

Exposure to various organic compounds including a number of environmental pollutants and drugs can cause cellular damage through generation of reactive oxygen species (ROS). Carbon tetrachloride (CCl<sub>4</sub>), a clear, colorless, volatile, heavy and nonflammable liquid, causes free radical generation and causes kidney injury in rats.<sup>[1,2]</sup> Free radicals induce lipid peroxidation and can damage cell membranes.<sup>[3]</sup> The average daily intake of CCl<sub>4</sub> for the general population is estimated to be 0.1 µg.

Kidney failure is frequently reported in fatal poisoning.<sup>[4,5]</sup> Furthermore, based on results of animal studies, the US Environmental Protection Agency has classified CCl<sub>4</sub> as a Group B2, probable human carcinogen.<sup>[5,6]</sup> In spite of the fact that the harmful effects of CCl<sub>4</sub> are obvious, this compound is still used as a solvent for oils, fats, lacquers, varnishes, rubber waxes and resins and as a starting material in the production of a number of organic compounds.<sup>[6,7]</sup>

It has been established that trichloromethyl (CCl<sub>3</sub>) radical and Cl are formed as a result of the metabolic conversion of CCl<sub>4</sub> by cytochrome P450, which in turn, initiate lipid peroxidation process.<sup>[8-10]</sup>

Among horticultural crops, fruits are sources of diverse nutrient and non-nutrient molecules, which display antioxidant properties,<sup>[11]</sup> and can protect the human body against oxidant damage.

*Cornus mas*, known as the European and Asiatic Cornelian Cherry, has been used for the treatment of diarrhea, intestinal inflammation, fever and malaria.<sup>[12]</sup> Furthermore, it has been mentioned for the treatment of kidney stones,

### Address for correspondence:

Miss. N. Banihabib, Drug Applied Research Center, Tabriz University of Medical Sciences, Golgasht Ave. Postal Code 5157658581, Tabriz-Iran. E-mail: n.banihabib@gmail.com

### Access this article online

#### Quick Response Code:



#### Website:

[www.indianjephrol.org](http://www.indianjephrol.org)

#### DOI:

10.4103/0971-4065.133000

kidney treatment and bladder infections in traditional system of medicine in Iran. Chemical characterization of *C. mas* fruit has shown that it is a rich source of phenolic and antioxidant, anthocyanins and flavonoids compounds.<sup>[12,13]</sup> Despite the favorable ethnopharmacological properties, its protective effect against CCl<sub>4</sub> nephrotoxicity has not been explored. In the present study, we investigated the effects of *C. mas* fruit on oxidative stress parameters in CCl<sub>4</sub>-induced nephrotoxicity in rats.

## Materials and Methods

### Chemicals

Trichloroacetic acid (TCA) and ethylenediaminetetraacetic acid (EDTA) were obtained from Sigma-Aldrich Chemical Co. Ltd. (USA). Thiobarbituric acid (TBA) and CCl<sub>4</sub> were obtained from Merck Co. (Germany). Assay kits for the estimation of creatinine, urea and acid uric were purchased from Pars Azma (Iran) and all other chemicals used were of analytical grade were obtained from either Sigma-Aldrich or Merck (Germany).

### Plant material

*C. mas* plant and its fruits were authenticated by the Botany Department of Tabriz University, Iran and obtained from suburbs of Arasbaran protected jungle (East Azerbaijan, Iran) at the end of spring 2012. The fruits were air-dried, protected from direct sunlight and powdered. The powder was kept in a closed container at 8°C.

### Extraction

A total of 500 g of powder was extracted with a mixture of methanol: water (7:3) at 25°C. The solvent was completely removed by rotary vacuum evaporator at 50°C. *C. mas* fruit extract (CMFE) was frozen at -20°C until use. The yield of the extract was 50% with reference to dry starting material.

### Toxicity study

For toxicity studies, groups of 10 mice were administered (i.g.) the test compounds in the doses 100-1650 mg/kg. The LD<sub>50</sub> (LD<sub>50</sub> = 1270) was determined using the graphical methods of Litchfield and Wilcoxon.<sup>[14]</sup> Two different doses were selected to evaluate the dose dependent effect of CMFE on CCl<sub>4</sub>-induced nephrotoxicity.

Furthermore, based on previous studies about evaluation of toxicity effects of CCl<sub>4</sub> like Zargar (2010), desired dose of CCl<sub>4</sub> was selected.

### Animals and treatment

Male albino rats of Wistar strain (250-300 g) were purchased from Pasteur Institute (Tehran, Iran). The animals were housed in polypropylene cages in a

temperature-controlled room (22 ± 2°C) with relative humidity (44-55%) under 12/12 h light and dark cycles for 1 week before and during the experiments. Animals were provided with a standard rodent pellet diet and clean drinking water *ad libitum*. Animals were divided into seven groups of six animals each:

- Group I served as a sham for both prophylactic and curative studies and received raw water and free access to food for 16 days
- Group II served as a normal control for both prophylactic and curative studies and received distilled water for 16 days orally and on the 16<sup>th</sup> day received olive oil (1 ml/kg body weight; i.p.)
- Group III served as a toxic control for both prophylactic and curative studies and received distilled water for 16 days orally and on the 16<sup>th</sup> day received CCl<sub>4</sub> (1 ml/kg body weight; 80% in olive oil)
- Groups IV and V served as pre-treatment groups (prophylactic). They received CMFE at doses of 300 and 700 mg/kg, orally for 16 days respectively and on the 16<sup>th</sup> day received CCl<sub>4</sub> (1 ml/kg body weight; 80% in olive oil), 2 h after administration of the last dose of extract
- Groups VI and VII served as post-treatment groups (curative). They received distilled water orally for 16 days and on the 16<sup>th</sup> day they received CCl<sub>4</sub> (1 ml/kg body weight; 80% in olive oil), followed by CMFE at doses of 300 mg/kg and 700 mg/kg (orally) respectively to Groups VI and VII at 2, 6, 12, 24 and 48 h after CCl<sub>4</sub> intoxication.

### Assessment of renal functions

All animals were sacrificed 50 h after CCl<sub>4</sub> administration. Blood samples were collected from left ventricle. Serum was separated by centrifugation at 3000 rpm for 15 min and used for biochemical estimations and was used freshly for the assessment of kidney function tests. The urea, acid uric, creatinine and total protein levels were estimated by standard diagnostic test kits (Pars Azma, Iran).

### Preparation of kidney homogenate

Renal tissues were homogenized in KCl (10 mM) phosphate buffer (1.15%) with EDTA: pH 7.4 and centrifuged at 12,000 rpm for 20 min. The supernatant was used for the measurement of malondialdehyde (MDA), catalase (CAT), superoxide dismutase (SOD) and glutathione peroxidase (GPx). Total protein contents were determined by the method of Lowry *et al.* (1951),<sup>[15]</sup> using bovine serum albumin as a standard.

### Measurement of lipid peroxidation

Lipid peroxidation was measured by the TBA reaction method.<sup>[16]</sup> In brief, samples were mixed with TBA reagent consisting of 0.375% TBA and 15% TCA in

0.25-N hydrochloric acid (HCl). The reaction mixtures were placed in a boiling water bath for 30 min and centrifuged at 2500 rpm for 5 min. The absorbance of the supernatant was measured at 535 nm. MDA, a measure of lipid peroxidation, was calculated using an extinction coefficient of  $1.56 \times 10^5/\text{M cm}$ . The results were expressed as nmol/mg protein.

#### Determination of antioxidant enzymes

CAT activity was measured according to the method of Aebi (1984).<sup>[17]</sup> One unit of CAT was defined as the amount of enzyme required to decompose 1  $\mu\text{M}$  of  $\text{H}_2\text{O}_2$  in 2 min. The reaction was initiated by the addition of 1.0 ml of freshly prepared 20 mM  $\text{H}_2\text{O}_2$ . The rate of decomposition of  $\text{H}_2\text{O}_2$  was measured spectrophotometrically at 240 nm for 1 min. The enzyme activity was expressed as U/mg protein.

The activity of SOD was measured according to the method of McCord *et al.* (1994).<sup>[18]</sup> For the determination of SOD activity, xanthine and xanthine oxidase were used to generate superoxide radicals reacting with 2-(4-iodophenyl)-3-(4-nitrophenol)-5-phenyl tetrazolium chloride to form a red formazan dye. SOD activity was measured at 505 nm. Data were expressed as U/mg of protein.

GPx activity was determined by the method described by Paglia and Valentine (1967).<sup>[19]</sup> The reaction mixture consisted of 400  $\mu\text{l}$ , 0.25 M potassium phosphate buffer (pH 7.0), 200  $\mu\text{l}$  supernatant, 100  $\mu\text{l}$  GSH (10 mM), 100  $\mu\text{l}$  NADPH (2.5 mM) and 100  $\mu\text{l}$  glutathione reductase (6 U/ml). Reaction was started by adding 100  $\mu\text{l}$  hydrogen peroxide (12 mM) and absorbance measured at 366 nm at 1 min intervals for 5 min using a molar extinction coefficient of  $6.22 \times 10^3/\text{M cm}$ . Data were expressed as U/mg of protein.

#### Histopathology

For microscopic evaluation kidneys were fixed in a fixative (absolute alcohol 60%, formaldehyde 30% and glacial acetic acid 10%) and embedded in paraffin, sectioned at 4  $\mu\text{m}$ , stained with hematoxylin/eosin and observed under a light microscope.

#### Animals rights and ethics

The study was cleared by the Institutional Animal Ethical Committee of Liver and Gastrointestinal Diseases Research Center (LGDR) (No 91/232 – LGDR).

#### Statistical analysis

All results are expressed as mean  $\pm$  standard error of the mean one way analysis of variance followed by multiple comparison. Chi-square test with the Tukey *post-hoc* and independent samples *T*-test was used to compared different parameters.  $P < 0.05$  was considered to be significant.

#### Results

##### Effect of CMFE on serum profile in rat

The effect of  $\text{CCl}_4$  administration on serum concentration of urea, acid uric, creatinine, protein and albumin are presented in Table 1. Administration of  $\text{CCl}_4$  to rats significantly ( $P < 0.05$ ) increased the level of creatinine, urea and uric acid while the protein and albumin levels were reduced in comparison with the control group. Treatment of CMFE significantly ( $P < 0.05$ ) reversed the changed levels of the above markers.

##### Effect of CMFE on renal antioxidant enzymes activity

Antioxidant enzymes activity in  $\text{CCl}_4$  group was found to be lower than in the normal group. The activity of these enzymes in treatment groups (pre and post) were significantly ( $P < 0.05$ ) increased when compared with  $\text{CCl}_4$  group that shown in Table 2.

##### Effect of CMFE on renal contents of MDA

Effect of  $\text{CCl}_4$  and CMFE on renal MDA is shown in Figure 1. Administration of  $\text{CCl}_4$  to rats significantly ( $P < 0.05$ ) induced lipid peroxidation as evidenced by the increased level of MDA. Treatment with CMFE protected against the  $\text{CCl}_4$  induced oxidative stress by reducing the lipid peroxidation.

##### Renal histopathology

Administration of  $\text{CCl}_4$  caused glomerular and tubular injury. Treatment with CMFE ameliorated the changes to near normal histology [Figure 2].

**Table 1: Effect of CMFE on serum profile in rat**

Treatment	Urea (mg/dl)	Creatinine (mg/dl)	Uric acid (mg/dl)	Total Protein (g/dl)	Albumin (g/dl)
Control	50.66 $\pm$ 1.11	0.89 $\pm$ 0.06	1.13 $\pm$ 0.13	7.85 $\pm$ 0.15	3.65 $\pm$ 0.06
Olive oil	59.66 $\pm$ 2.23*	1.07 $\pm$ 0.04*	1.35 $\pm$ 0.07*	7.93 $\pm$ 0.18*	3.38 $\pm$ 0.07*
1 ml/kg $\text{CCl}_4$	1.50 $\pm$ 3.76*	1.65 $\pm$ 0.04*	3.61 $\pm$ 0.16*	5.83 $\pm$ 0.23*	1.31 $\pm$ 0.08*
300 mg/kg CMFE+ $\text{CCl}_4$	84.50 $\pm$ 3.60*	1.24 $\pm$ 0.04*	1.88 $\pm$ 0.22*	7.10 $\pm$ 0.11*	2.20 $\pm$ 0.09*
700 mg/kg CMFE+ $\text{CCl}_4$	94.33 $\pm$ 3.06*	1.09 $\pm$ 0.03*	1.88 $\pm$ 0.17*	7.25 $\pm$ 0.14*	3.10 $\pm$ 0.09*
$\text{CCl}_4$ +300 mg/kg CMFE	67.50 $\pm$ 2.99*	1.33 $\pm$ 0.05*	1.78 $\pm$ 0.14*	7.71 $\pm$ 0.15*	2.85 $\pm$ 0.12*
$\text{CCl}_4$ +700 mg/kg CMFE	69.83 $\pm$ 4.04*	1.02 $\pm$ 0.05*	1.76 $\pm$ 0.13*	7.91 $\pm$ 0.16*	3.20 $\pm$ 0.09*

Results are expressed as mean $\pm$ SE. (n=6). \*Indicate significance at  $P < 0.05$  probability from control group. †Indicate significance at  $P < 0.05$  probability from  $\text{CCl}_4$  group. CMFE: *Cornus mas* fruit extract

## Discussion

The present study demonstrated the protective potential of CMFE on CCl<sub>4</sub> induced nephrotoxicity. CCl<sub>4</sub> has been used in rat experimental models to investigate the oxidative stress induced in various organs. To the best of our knowledge, this is the first study to evaluate these effects of CMFE in an attempt to prevent kidney damage from CCl<sub>4</sub>.

The mechanism of CCl<sub>4</sub> hepatotoxicity is well documented in the rat model.<sup>[14,20,21]</sup> CCl<sub>4</sub> intoxication generates free radicals that trigger a cascade of events resulting in organ toxicity in rats. It is well-known that the kidneys play a pivotal role in the regulation of various chemicals. CCl<sub>4</sub>, a nephrotoxin, was used for the purpose of inducing renal damage in this study since it has previously been shown to exert its toxic effects on the kidney.<sup>[22,23]</sup> According to previous reports, CCl<sub>4</sub>-induced toxicity is due to the conversion of CCl<sub>4</sub> to CCl<sub>3</sub>· and CCl<sub>3</sub>O<sub>2</sub>· by the liver cytochrome P450 enzyme. These highly reactive free radicals cause cell damage.

Elevations in the serum concentrations of urea and creatinine as seen here are indicative of renal injury,


which was accompanied by histological changes.<sup>[3,23,24]</sup> In addition, reduced level of serum albumin and protein in CCl<sub>4</sub>-treated rats might have resulted from leakage in glomeruli and tubules. Similar results have been shown by Khan *et al.* The present study revealed that the treatment of CMFE to CCl<sub>4</sub>-administrated rats ameliorated the toxic affect of CCl<sub>4</sub>. Results obtained in this study are in agreement with earlier findings.<sup>[25]</sup>

The lipid peroxidation is an autocatalytic process and common consequence of cell death. It causes tissue damage during inflammation, cancer and aging.<sup>[26]</sup> Lipid peroxidation is reported to be a major causes of CCl<sub>4</sub>-induced nephrotoxicity, mediated by the production of free radical derivatives of CCl<sub>4</sub>. The renal MDA content, which is one of the end products of lipid peroxidation in the renal tissue, is used as an important indicator of CCl<sub>4</sub>-induced oxidative stress. In the present study,


**Table 2: Effect of CMFE on renal antioxidant enzymes activity**

Treatment	U/mg protein		
	SOD	CAT	GPx
Control	30.69±0.77	10.91±0.49	62.19±1.64
Olive oil	29.73±0.77 <sup>+</sup>	9.91±0.34 <sup>+</sup>	58.25±1.79 <sup>+</sup>
1 ml/kg CCl <sub>4</sub>	19.54±0.39 <sup>*</sup>	4.46±0.24 <sup>*</sup>	27.33±1.74 <sup>*</sup>
300 mg/kg CMFE+CCl <sub>4</sub>	22.35±0.47 <sup>+</sup>	8.90±0.33 <sup>+</sup>	39.20±1.10 <sup>+</sup>
700 mg/kg CMFE+CCl <sub>4</sub>	23.22±0.35 <sup>+</sup>	9.60±0.22 <sup>+</sup>	42.36±0.95 <sup>+</sup>
CCl <sub>4</sub> +300 mg/kg CMFE	25.61±0.36 <sup>+</sup>	10.35±0.45 <sup>+</sup>	51.67±1.64 <sup>+</sup>
CCl <sub>4</sub> +700 mg/kg CMFE	24.30±0.36 <sup>+</sup>	9.49±0.23 <sup>+</sup>	47.14±1.64 <sup>+</sup>

Results are expressed as mean±SE. (n=6). <sup>\*</sup>Indicate significance at P<0.05 probability from control group. <sup>+</sup>Indicate significance at P<0.05 probability from CCl<sub>4</sub> group. CMFE: Cornus mas fruit extract



**Figure 1: Effect of CMFE on renal contents of MDA in rat.** <sup>\*</sup>Indicate significance at P < 0.05 probability from the control group. <sup>+</sup>Indicate significance at P < 0.05 probability from CCl<sub>4</sub> group. CMFE, Cornus mas fruit extract; MDA, malondialdehyde



**Figure 2: H and E stain:** (a) Representative section from the control group. (b) CCl<sub>4</sub> (1 ml/kg) group. (c) Olive oil (1 ml/kg) group. (d) CMFE (300 mg/kg) + CCl<sub>4</sub> group. (e) CMFE (700 mg/kg) + CCl<sub>4</sub> group. (f) CCl<sub>4</sub> + CMFE (300 mg/kg) group. (g) CCl<sub>4</sub> + CMFE (700 mg/kg) group. CMFE, Cornus mas fruit extract; CCl<sub>4</sub>, carbon tetrachloride; DBC, damaged Bowman capsule; TD, tubular degeneration; LBB, loss of border brush; PC, proximal tubule; DT, distal tubule; NGB, normal glomerulus and Bowman capsule; BB, border brush; FA, foamy appearance

administration of CCl<sub>4</sub> resulted in significant elevation in MDA concentration [Figure 1] indicating elevation of lipid peroxidation along with histopathological injury [Figure 2]. Interestingly, treatment by CMFE markedly the MDA concentration.

Although there are numerous studies demonstrating that CCl<sub>4</sub> leads to increase in MDA levels in various tissues,<sup>[27,28]</sup> a limited number of studies have investigated the effects of CMFE. In addition, no published data has ever demonstrated the influence of CMFE on CCl<sub>4</sub>-induced elevation in renal lipid peroxide levels.

It has been suggested that a decrease in the activities of primary antioxidant; CAT, SOD and GPx may be due accumulation of ROS. An observation that strengthens this hypothesis is that SOD activity can be inhibited by hydrogen peroxide treatment.<sup>[29]</sup> The inhibition of antioxidant system may lead to accumulation of H<sub>2</sub>O<sub>2</sub> or products of its decomposition may also be aided by a decrease in CAT, SOD and GPx activities.<sup>[2,30]</sup> Measurement of these antioxidant enzymes is an appropriate indirect way to assess the pro-oxidant antioxidant status in tissues.<sup>[30,31]</sup> The level of antioxidant enzymes such as SOD, CAT and GPx decreased in CCl<sub>4</sub>-treated group, and improved by treatment with CMFE. Results obtained in this study suggest the protective effects of CMFE against the CCl<sub>4</sub>-induced nephrotoxicity, could be attributed to its high level of phenol<sup>[32]</sup> and other antioxidants.<sup>[33-37]</sup> These compounds could scavenge the free radicals of CCl<sub>4</sub> generated through P450 enzyme system thereby diminished the oxidative injuries.

In this study, the kidneys of CCl<sub>4</sub>-treated rats have shown severe morphological abnormalities in the glomerular and tubular compartments. These changes were not observed in the groups treated with CMFE, that suggesting the protective effects of CMFE in attenuating CCl<sub>4</sub>-induced morphological changes.

## Conclusion

The present study suggests the antioxidant potential of CMFE against the toxic effects of CCl<sub>4</sub> in the kidney of rats. Research is needed about each of these components against CCl<sub>4</sub> induced nephrotoxicity.

## References

- Khan MR, Rizvi W, Khan GN, Khan RA, Shaheen S. Carbon tetrachloride-induced nephrotoxicity in rats: Protective role of *Digera muricata*. J Ethnopharmacol 2009;122:91-9.
- Khan RA, Khan MR, Sahreen S. Evaluation of *Launaea procumbens* use in renal disorders: A rat model. J Ethnopharmacol 2010;128:452-61.
- Ogeturk M, Kus I, Colakoglu N, Zararsiz I, Ilhan N, Sarsilmaz M. Caffeic acid phenethyl ester protects kidneys against carbon tetrachloride toxicity in rats. J Ethnopharmacol 2005;97:273-80.
- Faroon O, Taylor J, Roney N, Fransen ME, Bogaczyk S, Diamon G. Toxicological profile for carbon tetrachloride: Agency for Toxic Substances and Disease Registry 2005.
- Soderland P, Lovekar S, Weiner DE, Brooks DR, Kaufman JS. Chronic kidney disease associated with environmental toxins and exposures. Adv Chronic Kidney Dis 2010;17:254-64.
- Thrall KD, Vucelick ME, Gies RA, Zangar RC, Weitz KK, Poet TS, et al. Comparative metabolism of carbon tetrachloride in rats, mice, and hamsters using gas uptake and PBPK modeling. J Toxicol Environ Health A 2000;60:531-48.
- Abraham P, Wilfred G, Cathrine. Oxidative damage to the lipids and proteins of the lungs, testis and kidney of rats during carbon tetrachloride intoxication. Clin Chim Acta 1999;289:177-9.
- Adewole S, Salako A, Doherty O, Naicker T. Effect of melatonin on carbon tetrachloride-induced kidney injury in Wistar rats. Afr J Biomed Res 2010;10:153-164.
- Yuan LP, Chen FH, Ling L, Bo H, Chen ZW, Li F, et al. Protective effects of total flavonoids of *Bidens bipinnata* L. against carbon tetrachloride-induced liver fibrosis in rats. J Pharm Pharmacol 2008;60:1393-402.
- Ugur H, Amat N, Blazeković B, Talip A. Protective effect of *Cichorium glandulosum* root extract on carbon tetrachloride-induced and galactosamine-induced hepatotoxicity in mice. Food Chem Toxicol 2009;47:2022-30.
- Scalzo J, Politi A, Pellegrini N, Mezzetti B, Battino M. Plant genotype affects total antioxidant capacity and phenolic contents in fruit. Nutrition 2005;21:207-13.
- Zargari A. Medicinal plants. Tehran University Publications, Tehran 1996;513:38.
- Serteser A, Kargoglu M, Gök V, Bağcı Y, Özcan MM, Arslan D. Antioxidant properties of some plants growing wild in Turkey. Grasas y Aceites 2009;60:147-54.
- Sundari PN, Wilfred G, Ramakrishna B. Does oxidative protein damage play a role in the pathogenesis of carbon tetrachloride-induced liver injury in the rat? Biochim Biophys Acta 1997;1362:169-76.
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the Folin phenol reagent. J Biol Chem 1951;193:265-75.
- Berton TR, Conti CJ, Mitchell DL, Aldaz CM, Lubet RA, Fischer SM. The effect of vitamin E acetate on ultraviolet-induced mouse skin carcinogenesis. Mol Carcinog 1998;23:175-84.
- Aebi H. Catalase *in vitro*. Methods in Enzymology. Academic Press 1984;105:121-26.
- McCord JM. Mutant mice, Cu, Zn superoxide dismutase, and motor neuron degeneration. Science 1994;266:1586-7.
- Paglia DE, Valentine WN. Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. J Lab Clin Med 1967;70:158-69.
- Sharma M, Pillai K, Husain S, Giri D. Protective role of propolis against alcohol carbon tetrachloride-induced hepatotoxicity in rats. Indian J Pharmacol 1997;29:76.
- Sotelo-Félix JI, Martínez-Fong D, Muriel P, Santillán RL, Castillo D, Yahuaca P. Evaluation of the effectiveness of *Rosmarinus officinalis* (Lamiaceae) in the alleviation of carbon tetrachloride-induced acute hepatotoxicity in the rat. J Ethnopharmacol 2002;81:145-54.
- Doğukan A, Akpolat N, Celiker H, Ilhan N, Halil Bahçecioğlu I, Günal AI. Protective effect of interferon-alpha on carbon tetrachloride-induced nephrotoxicity. J Nephrol 2003;16:81-4.
- Ozturk F, Ucar M, Ozturk IC, Vardi N, Batcioglu K. Carbon tetrachloride-induced nephrotoxicity and protective effect of betaine in Sprague-Dawley rats. Urology 2003;62:353-6.
- Adelman RD, Spangler WL, Beasom F, Ishizaki G, Conzelman GM.

- Furosemide enhancement of experimental gentamicin nephrotoxicity: Comparison of functional and morphological changes with activities of urinary enzymes. *J Infect Dis* 1979;140:342-52.
25. Shyu MH, Kao TC, Yen GC. Hsian-tsao (*Mesona procumbens* Heml.) prevents against rat liver fibrosis induced by CCl<sub>4</sub> (4) via inhibition of hepatic stellate cells activation. *Food Chem Toxicol* 2008;46:3707-13.
  26. Bhadauria M, Nirala SK. Reversal of acetaminophen induced subchronic hepatorenal injury by propolis extract in rats. *Environ Toxicol Pharmacol* 2009;27:17-25.
  27. Daniels WM, Reiter RJ, Melchiorri D, Sewerynek E, Pablos MI, Ortiz GG. Melatonin counteracts lipid peroxidation induced by carbon tetrachloride but does not restore glucose-6 phosphatase activity. *J Pineal Res* 1995;19:1-6.
  28. Donder E, Baydas G, Ozkan Y, Ercel E, Yalniz M, Dogan H. Investigation of antioxidant effect of melatonin against carbon tetrachloride toxicity in various tissues. *Biomed Res* 1999;10:141-5.
  29. Miguel F, Augusto AC, Gurgueira SA. Effect of acute vs chronic H<sub>2</sub>O<sub>2</sub>-induced oxidative stress on antioxidant enzyme activities. *Free Radic Res* 2009;43:340-7.
  30. Halliwell B. Reactive oxygen species in pathology with special reference to the skin. *Oxidative Stress in Dermatology*. New York: Marcek Dekker Inc.; 1993.
  31. Priscilla DH, Prince PS. Cardioprotective effect of gallic acid on cardiac troponin-T, cardiac marker enzymes, lipid peroxidation products and antioxidants in experimentally induced myocardial infarction in Wistar rats. *Chem Biol Interact* 2009;179:118-24.
  32. Afolayan AJ, Jimoh FO. Nutritional quality of some wild leafy vegetables in South Africa. *Int J Food Sci Nutr* 2009;60:424-31.
  33. Shimizu S, Miyase T, Ueno A, Usmanghani K. Sesquiterpene lactone glycosides and ionone derivative glycosides from *Sonchus asper*. *Phytochemistry* 1989;28:3399-402.
  34. Giner RM, Ubeda A, Just MJ, Serrano A, Máñez S, Ríos JL. A chemotaxonomic survey of *Sonchus* subgenus *Sonchus*. *Biochem Syst Ecol* 1993;21:617-20.
  35. Máñez S, Recio MC, Giner RM, Sanz MJ, Terencio MC, Peris JB, et al. A chemotaxonomic review of the subtribe crepidinase based on its phenol constituents. *Biochem Syst Ecol* 1994;22:297-305.
  36. Guil Guerrero JL, Giménez Giménez A, Rodríguez García I, Torija Isasa ME. Nutritional composition of *Sonchus* species (*S. asper* L., *S. oleraceus* L. and *S. tenerimus* L.). *J Sci Food Agric* 1998;76:628-32.
  37. Helal AM, Nakamura N, El-Askary H, Hattori M. Sesquiterpene lactone glucosides from *Sonchus asper*. *Phytochemistry* 2000;53:473-7.
- How to cite this article:** Es.Haghi M, Dehghan G, Banihabib N, Zare S, Mikaili P, Panahi F. Protective effects of *Cornus mas* fruit extract on carbon tetrachloride induced nephrotoxicity in rats. *Indian J Nephrol* 2014;24:291-6.

**Source of Support:** Nil, **Conflict of Interest:** None declared.

